

## ENTOMOLOGICAL NEWS, THE AMERICAN ENTOMOLOGICAL SOCIETY, AND NEW GUIDELINES FOR AUTHORS OF ENTOMOLOGICAL NEWS

*Entomological News* is published bimonthly except July-August by The American Entomological Society, which is headquartered at the Academy of Natural Sciences, 1900 Benjamin Franklin Parkway, Philadelphia, PA 19103-1195 United States of America. AES can be reached, as follows: telephone (215) 561-3978; fax (215) 299-1028; e-mail, aes@acnatsci.org and website: <http://www.acnatsci.org/hosted/aes>.

**Subscriptions to and Back Issues of *Entomological News*.** Private subscriptions for personal use of members of the society are US \$15 per year prepaid. Subscriptions for institutions, such as libraries, laboratories, etc. and nonmembers, are US\$30 per year prepaid for those located in the USA and US \$34 per year prepaid for those located outside the USA. Back issues when available are sold by complete volume, for US \$15 to members, and US \$30 to nonmembers. Membership / subscription application and additional information is available at: <http://www.acnatsci.org/hosted/aes/subscription.html>. Please send inquiries or send completed membership form to: Office Manager at the address above, e-mail: aes@say.acnatsci.org, or call (215) 561-3978.

**Previous editors of *Entomological News*:** 1890 Eugene Murray-Aaron (1852-1940); 1890-1920 Henry Skinner (1861-1926); 1921-1944 Philip P. Calvert (1871-1961); 1945-1967 R.G. Schmieder (1898-1967); 1968-1972 Ross H. Arnett, Jr. (1919-1999); 1973-1974 R.W. Lake; 1974-2002 Howard P. Boyd; 2002-2003 F. Christian Thompson and Michael Pogue.

### New Guidelines for authors of *Entomological News*:

Further guidelines can be found on <http://www.geocities.com/entomologicalnews/instructions.htm>

**Subject Coverage:** *Entomology, sensu lato*. Manuscripts on systematics, ecology, evolution, morphology, physiology, behavior, biodiversity, conservation, paleobiology, and other aspects of insect and terrestrial arthropod life as well as nomenclature, biographies and history of entomology, among others, are appropriate topics for papers submitted to *Entomological News*. Papers on applied, economic, and regulatory entomology or on toxicology and related subjects will be considered only if they also make a major contribution in one of the aforementioned fields.

**Any author may submit papers.** Manuscripts will be accepted from any author, although, papers from members of the American Entomological Society are given priority. It is suggested that all prospective authors join the AES.

**Send manuscripts, books for review, and editorial correspondence to the editor.** All manuscripts, including scientific notes and book reviews, submitted for publication in *Entomological News* as well as all associated editorial communications must be sent to the Editor, Jorge A. Santiago-Blay at this address: Department of Paleobiology, MRC-121, National Museum of Natural History, Smithsonian Institution, P. O. Box 37012, Washington, D.C. 20013-7012 U.S.A. If an author uses a mailing service that does not accept addresses with a P. O. Box, please use this address: Department of Paleobiology, MRC-121, National Museum of Natural History, Smithsonian Institution, West Loading Dock, Washington, District of Columbia 20560 U.S.A. Other means of contacting the Editor are, as follows: Phone: (202) 633-1383, Fax: (202) 786-2832, e-mails: [santiago-blay@nmnh.si.edu](mailto:santiago-blay@nmnh.si.edu), or [blayj@hotmail.com](mailto:blayj@hotmail.com), or via the webpage at <http://www.geocities.com/entomologicalnews/contact.htm>. Books for review should also be sent to the editor who will, in turn, give them to a colleague for review.

**Agreement.** By submitting a paper for publication, authors tacitly agree to not submit in parallel the same manuscript to another journal. For published papers, authors agree to accept responsibility for all page, illustration, and requested reprint charges. Rejected manuscripts will be discarded, except for original artwork and photographs, which will be returned to the authors.

**Scientific Notes and Book Reviews.** These are much shorter contributions, typically not exceeding two printed pages. The main difference between these types of contributions and longer papers is that scientific notes and book reviews lack an abstract and most of the main headings, except for the acknowledgments and the literature cited sections.

Postmaster: If undeliverable, please send to

The American Entomological Society at The Academy of Natural Sciences  
1900 Benjamin Franklin Parkway / Philadelphia, PA 19103-1195

THIRD CLASS POSTAGE PAID AT DOVER, DELAWARE 19901 U.S.A.

## TAXONOMIC STATUS OF THE AUSTRALIAN MAYFLY GENERA *JAPPA* AND *ULMEROPHLEBIA* (EPHEMEROPTERA: LEPTOPHLEBIDAE)<sup>1</sup>

Y. J. Bae,<sup>2</sup> K. J. Finlay,<sup>3</sup> and I. C. Campbell<sup>1</sup>

**ABSTRACT:** The Australian burrowing mayfly genera *Jappa* and *Ulmerophlebia* are confirmed as monophyletic groups. Their adult and larval stages are redescribed and their egg stages are newly described. Additional taxonomic and phylogenetic discussions are provided.

**KEY WORDS:** Ephemeroptera, Leptophlebiidae, *Jappa*, *Ulmerophlebia*, Australia.

The Australian mayfly genera *Jappa* Harker and *Ulmerophlebia* Demoulin (Leptophlebiidae) are unique among Ephemeroptera because of the convergent adaptation with the burrowing mayflies (Ephemeroidea) from the Northern Hemisphere (Campbell, 1990; Bae and McCafferty, 1991, 1995). Members of the Ephemeroidea do not occur in Australia, and these are the genera of Ephemeroptera most closely adapted to hyporheic habitats in Australia. The larvae are generally found under cobble- or boulder-sized stones embedded in sand and silt substrate in mid-sized to large lowland streams (Riek, 1970; Peters and Campbell, 1991; Edmunds and McCafferty, 1996).

The larvae of *Jappa* possess a characteristic head with a two-pronged frontal process or "cephalic tusks" (Fig. 1) that is analogous to the mandibular tusks of Ephemeroidea, in particular those of Potamanthidae (see Bae and McCafferty, 1991; Figs. 11-14), while the larvae of *Ulmerophlebia* lack such structure (Fig. 2). Despite this morphological difference, the generic distinction of the genera has been continuously questioned by mayfly taxonomists not only because their general morphology in adult and larval stages is similar, but because generic concepts have not been well defined.

Harker (1954) established the genus *Jappa* based on adult and larval stages. At that time, forewing length to width ratio and "a burrowing type larva" with tusk-like head frontal processes or "horns" were used as the major defining characteristics. Demoulin (1955) erected the genus *Ulmerophlebia* to include a species, *Euphyurus mjobergi* Ulmer, described (as an adult only) by Ulmer (1916), but did not compare it with *Jappa*. Previously, Ulmer (1920) recombined *E. mjobergi* with *Deleatidium* Eaton; and Demoulin (1955) distinguished *Ulmerophlebia* from *Deleatidium* mainly by wing and genital characters.

<sup>1</sup> Received on June 27, 2002. Accepted on May 18, 2004.

<sup>2</sup> Department of Biology, Seoul Women's University, Seoul 139-774, Korea. E-mail: ybae@swu.ac.kr.

<sup>3</sup> School of Biological Sciences, Monash University, Melbourne, Victoria 3800, Australia. E-mail: ian.campbell@sci.monash.edu.au.



Figs. 1-2. Larval head: 1. *Jappa kutera* (bar = 0.5mm). 2. *Ulmerophlebia mjobergi* (bar = 0.5mm).

Williams (1968) mentioned a possible congeneric status of *Jappa* and *Ulmerophlebia* based on personal communication with E. F. Riek. Riek (1970), in the Ephemeroptera chapter of the textbook "The Insects of Australia," placed all burrowing Australian leptophlebiids, i.e. *Jappa* and *Ulmerophlebia*, into *Jappa* without any explanations. Peters and Campbell (1991) also followed the previous classification by Riek (1970) in the second edition of the textbook. Suter (1986) provided a historical background of *Ulmerophlebia* and described the larval stage of *Ulmerophlebia* for the first time based on a second species, *U. pipinna* Suter, but was conservative in clarifying the generic status of the *Ulmerophlebia* and *Jappa*. Dean (1999) gave larval diagnoses of *Jappa* and *Ulmerophlebia* when he provided larval keys to three nominal and four unnamed species of *Jappa* and four unnamed species of *Ulmerophlebia*, but still did not resolve the generic status.

For the above reasons, the generic concepts of *Jappa* and *Ulmerophlebia* have not been fully resolved. The purpose of this study is to clarify and delineate the genera and provide detailed redescrptions of adult, larval, and egg stages.


Type and voucher specimens and additional fresh materials of all previously known species of *Jappa* and *Ulmerophlebia* (see Species included, pp. 5, 7) are housed mainly in the Museum of Victoria, Australian National Insect Collection in Canberra, and Monash University and were examined for this study. Terminology, measurement, and other general methods are after Bae and McCafferty (1991).

## *Jappa* Harker

*Jappa* Harker, 1954: 257 [Type species: *Jappa kutera* Harker, by original designation; M & L]; Williams, 1968: 170 (L key); Dean and Suter, 1996: 44 (L); Dean, 1999: 34 (L).

**Adult.** Male body length 9.4-11.3 mm; female body 12.0-16.5 mm; caudal filaments 14.0-19.1 mm. General body color light yellow with dark purplish brown markings. **Head:** Male dorsal compound eyes 1.10-1.43 mm in length, 1.00-1.19 mm in width, 0.48-0.68 mm in height, broadly meeting posteromedially (B/D = 0), dome-shaped, dorsally oval, and with anterior margin round and posterior margin somewhat attenuating from lateral view ("bicycle helmet" shaped); basal compound eyes anteroventrally pronounced, invisible to slightly visible from dorsal view, 0.67-0.87 mm in length, 0.43-0.58 mm in height, and anteriorly oriented from lateral view. Female compound eyes 0.33-0.47 mm in width and 1.53-1.67 mm in distance between compound eyes (B/D = 3.50-4.70). **Thorax:** Forewings transparent, often with dark purplish brown markings at basal, central, and apical areas, 8.8-15.6 mm in length, and 2.9-5.3 mm in width; longitudinal veins light brown to dark purplish brown; crossveins light brown to dark purplish brown (basally and anteriorly located crossveins often darker and more heavily infuscated); crossveins C-Sc 20-30, Sc-R1 16-20, and R1-R2 13-15 in number; crossveins in stigmatic area not anastomosed; MP2 basally connected to MP1 and CuA; ICu1 basally connected to CuA and CuP (angle between ICu1 and crossvein ICu1-CuP larger than angle between ICu1 and crossvein ICu1-CuA). Hindwings transparent, 2.3-2.9 mm in length, and 1.2-1.4 mm in width; longitudinal veins white to light yellow; crossveins C-Sc, Sc-R1, and some in distal part sometimes dark brown; Sc ending relatively distally (length of Sc / distance from base of Sc to apex of distal margin = 0.89); crossveins C-Sc 9-11, more or less evenly distributed (in one species, C-Sc crossveins apically concentrated); costal margin round, with weakly developed costal projection at midlength; Rs 0.48-0.68 mm; R1 0.64-0.98 mm; MP3 0.25-0.48 mm; MP1 1.11-1.37 mm. Male forefemora 2.10-2.27 mm, foretibiae 2.90-3.25 mm, foretarsal segments 1, 2, 3, 4, and 5 0.13-0.14 mm, 1.08-1.27 mm, 1.00-1.18 mm, 0.76-0.89 mm, and 0.38-0.44 mm (foretarsal segment 2-3 > 4 > 5 > 1), and foreclaws 0.15-0.24 mm. Female foretibiae > forefemora > foretarsi. Claws dissimilar. **Abdomen:** Abdominal segment 9 with moderately to well developed posterolateral projections. Penis Y-shaped or V-shaped, relatively long (height of penis / height of forceps segment 1 = 0.45-0.56); forceps 3-segmented, with segment 1 0.92-1.03 mm, segment 2 0.10-0.13 mm, and segment 3 0.08-0.12 mm in length; forceps segment 1 basally broad and apically slender, with relatively abrupt constriction (constriction angle 80-110°) at mid-length to 3/4 apically; forceps segments 2 and 3 often indistinctly demarcated. Caudal filaments light yellow, often with purplish brown to dark brown stripes, often with maculation at joints; cerci 1.3-1.7x length of body; terminal filament as long as cerci.

**Larva.** Male body length 10.4-16.9 mm; cephalic tusks 1.0-2.0 mm; caudal filaments 5.9-10.9 mm. Female body length 10.4-23.8 mm; cephalic tusks 1.4-3.5 mm; caudal filaments 5.9-18.1 mm. Body surface glassy and heavily setose. General body color light yellow to light brown with dark purplish brown markings. **Head** (Fig. 1) 1.30-2.38 mm in length and 1.75-4.13 mm in width. Male compound eyes dorsally 0.56-0.60 mm in width and 0.65-1.06 mm in distance between compound eyes (B/D = 1.08-1.76). Female compound eyes dorsally 0.23-0.56 mm in width and 1.38-2.86 mm in distance between compound eyes (B/D = 4.75-6.11). Antennae 4.4-6.3 mm in length, with whorls of hairlike setae at each segment. Cephalic tusks arched and attenuating, apically convergent and upward, 1.32-2.30 mm in inner length and 1.58-2.88 mm in outer length, 14.9-26.5° in curvature; each cephalic tusk with rudimentary to distinct dorsal ridge, with small to large basodorsal tubercle, with fields of 6-20, 5-15, and 10-20 hairlike setae at basomedial, basosubapical, and apicodorsal areas (in two species apicodorsal setal field lacking), respectively, sometimes with additional dorsolateral row of 6-13 spines. Labrum distally wider (maximum length 0.30-0.38 mm; basal width 0.59-0.87 mm; distal maximum width 0.75-1.06 mm), dorsal surface heavily setose, with basal and subapical hairlike setal fringes (basal setae longer than subapical setae); subapical setal fringe laterally longer and 8-15 setae densely arranged medially; anterior and lateral margins with hairlike setal row; anterior margin concave, with prominent median tubercle, and with three pairs of rudimentary submedian denicles; ventral surface with dense hairlike setal field along anterior margin, with row of 10-20 stout setae on subanterior margins, and with fields of 20-50 hairlike setae at submedian areas.




Figs. 1-2. Larval head: 1. *Jappa kutera* (bar = 0.5mm). 2. *Ulmerophlebia mjobergi* (bar = 0.5mm).

Williams (1968) mentioned a possible congeneric status of *Jappa* and *Ulmerophlebia* based on personal communication with E. F. Riek. Riek (1970), in the Ephemeroptera chapter of the textbook "The Insects of Australia," placed all burrowing Australian leptophlebiids, i.e. *Jappa* and *Ulmerophlebia*, into *Jappa* without any explanations. Peters and Campbell (1991) also followed the previous classification by Riek (1970) in the second edition of the textbook. Suter (1986) provided a historical background of *Ulmerophlebia* and described the larval stage of *Ulmerophlebia* for the first time based on a second species, *U. pipinna* Suter, but was conservative in clarifying the generic status of the *Ulmerophlebia* and *Jappa*. Dean (1999) gave larval diagnoses of *Jappa* and *Ulmerophlebia* when he provided larval keys to three nominal and four unnamed species of *Jappa* and four unnamed species of *Ulmerophlebia*, but still did not resolve the generic status.

For the above reasons, the generic concepts of *Jappa* and *Ulmerophlebia* have not been fully resolved. The purpose of this study is to clarify and delineate the genera and provide detailed redescriptions of adult, larval, and egg stages.

Type and voucher specimens and additional fresh materials of all previously known species of *Jappa* and *Ulmerophlebia* (see Species included, pp. 5, 7) are housed mainly in the Museum of Victoria, Australian National Insect Collection in Canberra, and Monash University and were examined for this study. Terminology, measurement, and other general methods are after Bae and McCafferty (1991).

Mandible dorsolateral margins with very long hairlike setal row; ventral surface with transverse row of 10-20 hairlike setae; inner incisors slightly smaller to as large as outer incisors; incisors with 2-3 apical teeth and 0-4 lateral denticles; protheca rudimentary, with well-developed fringe. Hypopharynx superlinguae laterally curved and apically pointed. Maxillae with dense hairlike setal field medially 3/4 on galeolacinal crown, mixed with rowed comblike setae medially 1/2 on galeolacinal crown, with one pronounced comblike seta medioapically, and with rowed dense hairlike setae on medial margin; maxillary palp segment 1 0.27-0.52 mm, segment 2 0.37-0.48 mm, and segment 3 0.19-0.37 mm in length; segment 1 and 2 with sparse hairlike setae alone outer margin; segment 3 indistinctly demarcated from segment 2, with pronounced outer margin, and with strongly developed setal field along outer margin. Labial glossae dorsoventrally elongated and ventrally stalked, with dense hairlike setae; paraglossae with dense hairlike setal field dorsoapically; labial palp segment 1 0.28-0.59 mm, segment 2 0.31-0.40 mm, and segment 3 0.22-0.32 mm in length; segment 3 indistinctly demarcated from segment 2, apically pointed, with dense hairlike setal field along outer margin, and with stout setal row along inner margin. *Thorax*: Pronotum anterolateral margins round; lateral margins with row of sparse to dense hairlike setae. Forefemora 1.67-4.88 mm, foretibiae 1.67-5.63 mm, foretarsi 0.71-1.63 mm, and foreclaws 0.24-0.63 mm in length (foretibiae>forefemora>foretarsi>foreclaws); forefemora with long hairlike setal fields along anterior and posterior margins and basomedial area on dorsal surface; foretibiae with dense hairlike setal field (filtering setae) rowed along inner and outer margins (dorsomedially bare), with stout setal field (raking setae) on 2/3 apically to entire inner margin; foretarsi with dense hairlike setae on dorsal and lateral surfaces; foreclaws apically darker, with 10-15 tiny teeth 3/4 basally. Midlegs and hindlegs heavily setose; length femora>tibiae>tarsi>claws. *Abdomen*: Terga light brown to light purplish brown, mostly with submedian and sublateral dark purplish brown stripes, with very long hairlike setae covered on 1/2-3/4 dorsal area along median line, and with hairlike setal row along lateral margins; abdominal segment 8-9 with moderately to well developed posterolateral projections. Sterna bare (sterna 9-10 sometimes with hairlike setae). Gills on abdominal segment 1-7, double; both lamellae with indistinct to distinct



Figs. 3-6. Eggs: 3. *Jappa kutera* (bar = 38µm). 4. *J. kutera*, in part (bar = 10µm). 5. *Umerophlebia mjobergi* (bar = 43µm). 6. *U. mjobergi*, in part (bar = 17.6µm).

tracheae, with single apical filament; gill lamella inner part strongly expanded apicolaterally, and with fine setae on 1/5-1/2 apical margin; gill lamella outer part with fine setae on entire margin; apical filament attenuating, marginally with fine setae; gills 4 1.38-2.86 mm in length, 0.78-1.83 mm in width, and 1.38-2.38 mm in filament length, with weakly to strongly developed apical expansion. Caudal filament segments with whorls of setae.

**Egg** (Fig. 3) oval; long axis 124 µm; short axis 80 mm. Color light yellow in nature, white in alcohol. Egg surface (Fig. 4) with ca. 780 knob-terminated coiled threads relatively evenly distributed throughout egg surface; diameter of knob-terminated coiled threads 3.9-4.5 mm. Polar caps absent. Micropyles several, scattered, tageniform; sperm guide circular, 2.6-3.3 µm in diameter.

**Diagnosis.** Adults of *Jappa* possess greater numbers of C-Sc crossveins (20-30) in the forewings and more or less evenly distributed C-Sc crossveins in the hindwings (excluding one species), while those of *Umerophlebia* possess fewer C-Sc crossveins (12-19) in the forewings and apically concentrated C-Sc crossveins in the hindwings. The hindwing vein Sc of *Jappa* ends relatively distally (length of Sc / distance from base of Sc to apex of distal margin = 0.89) comparing with that of *Umerophlebia* (0.76). The penes of *Jappa* are relatively longer (height of penis / height of forceps segment 1 = 0.45-0.56) than those of *Umerophlebia* (0.08-0.31). The constriction in the medial margin of forceps segment 1 of *Jappa* is relatively abrupt (angle 80-110°), while that of *Umerophlebia* is relatively gradual (angle 135-150°). The larvae of *Jappa* can easily be distinguished from those of *Umerophlebia* and any other leptophlebid genera by the cephalic tusks (Fig. 1). In addition, the body size of *Jappa* in both of the adults and larvae (adult 9.4-16.5 mm, larva 10.4-23.8 mm) is generally larger than that of *Umerophlebia* (adult 6.9-8.7 mm, larva 5.5-10.0 mm).

**Species included.** *Jappa kutera* Harker (1954), *J. edmundsi* Skedros and Polhemus (1986), *J. serrata* Skedros and Polhemus (1986), *Jappa* AV1 (Dean, 1999), *Jappa* AV2 (Dean, 1999), *Jappa* AV3 (Dean, 1999), and *Jappa* AV4 (Dean, 1999).


**Distribution.** NSW, NT, northern WA, QLD, and VIC.

**Remarks.** There are several other species previously considered as, or assigned to, *Jappa*. *Jappa tristis* Harker (1954) [Holotype stage: M; locality: Tasmania, Cradle Mt.; deposition: British Museum (Natural History)] was subsequently identified as *Tillyardophlebia* Dean by Dean (1999). *Jappa* is not considered present in Tasmania.

*Leptophlebia furcifera* Eaton (1871) [Type stage: M; locality: Melbourne; deposition: Melbourne Museum = Museum of Victoria], recombined with *Atalophlebia* Eaton by Eaton (1884), was considered in *Jappa* by Dean (1999). However, the type specimen was not preserved in the Museum of Victoria when YJB checked in 2001 and there is little evidence that this species belongs to *Jappa* based on Eaton's original description (Eaton, 1871) and redescription (Eaton, 1884).

*Leptophlebia strigata* Eaton (1871) [Type stage: F; locality: North Australia; deposition: McLachlan Collection in British Museum (Natural History)], recombined with *Atalophlebia* by Eaton (1884) and *Deleatidium* by Ulmer (1920), was also considered in *Jappa* by Dean (1999). The original description (Eaton, 1871) and redescription (Eaton, 1884) of the species do not clearly substantiate the

Mandible dorsolateral margins with very long hairlike setal row; ventral surface with transverse row of 10-20 hairlike setae; inner incisors slightly smaller to as large as outer incisors; incisors with 2-3 apical teeth and 0-4 lateral denticles; prostheca rudimentary, with well-developed fringe. Hypopharynx superlinguae laterally curved and apically pointed. Maxillae with dense hairlike setal field medially 3/4 on galealacinial crown, mixed with rowed comblike setae medially 1/2 on galealacinial crown, with one pronounced comblike seta medioapically, and with rowed dense hairlike setae on medial margin; maxillary palp segment 1 0.27-0.52 mm, segment 2 0.37-0.48 mm, and segment 3 0.19-0.37 mm in length; segment 1 and 2 with sparse hairlike setae along outer margin; segment 3 indistinctly demarcated from segment 2, with pronounced outer margin, and with strongly developed setal field along outer margin. Labial glossae dorsoventrally elongated and ventrally stalked, with dense hairlike setae; paraglossae with dense hairlike setal field dorsoapically; labial palp segment 1 0.28-0.59 mm, segment 2 0.31-0.40 mm, and segment 3 0.22-0.32 mm in length; segment 3 indistinctly demarcated from segment 2, apically pointed, with dense hairlike setal field along outer margin, and with stout setal row along inner margin. *Thorax*: Pronotum anterolateral margins round; lateral margins with row of sparse to dense hairlike setae. Forefemora 1.67-4.88 mm, foretibiae 1.67-5.63 mm, foretarsi 0.71-1.63 mm, and foreclaws 0.24-0.63 mm in length (foretibiae  $\geq$  forefemora > foretarsi > foreclaws); forefemora with long hairlike setal fields along anterior and posterior margins and basomedial area on dorsal surface; foretibiae with dense hairlike setal field (filtering setae) rowed along inner and outer margins (dorsomedially bare), with stout setal field (raking setae) on 2/3 apical to entire inner margin; foretarsi with dense hairlike setae on dorsal and lateral surfaces; foreclaws apically darker, with 10-15 tiny teeth 3/4 basally. Midlegs and hindlegs heavily setose; length femora > tibiae > tarsi > claws. *Abdomen*: Terga light brown to light purplish brown, mostly with submedian and sublateral dark purplish brown stripes, with very long hairlike setae covered on 1/2-3/4 dorsal area along median line, and with hairlike setal row along lateral margins; abdominal segment 8-9 with moderately to well developed posterolateral projections. Sterna bare (sterna 9-10 sometimes with hairlike setae). Gills on abdominal segment 1-7, double; both lamellae with indistinct to distinct



Figs. 3-6. Eggs: 3. *Jappa kutera* (bar = 38 $\mu$ m). 4. *J. kutera*, in part (bar = 10 $\mu$ m). 5. *Ulmerophlebia mjobergi* (bar = 43 $\mu$ m). 6. *U. mjobergi*, in part (bar = 17.6 $\mu$ m).



generic position, and it is crucial that the types be reexamined before generic placement can be confirmed.

### *Ulmmerophlebia* Demoulin

*Ulmmerophlebia* Demoulin, 1955: 228 [Type species: *Euphyurus mjobergi* Ulmer, by original designation; A]; Suter, 1986: 352 [M & L]; Dean and Suter, 1996: 44 [L]; Dean, 1999: 74 [L].

**Adult.** Male body length 6.9–8.7 mm; female body 6.3–8.6 mm; caudal filaments 7.5–9.5 mm. General body color light yellow with dark purplish brown markings. **Head:** Male dorsal compound eyes 0.92–0.98 mm in length, 0.78–0.87 mm in width, 0.44–0.56 mm in height, 0–0.05 mm in distance between compound eyes (B/D = 0–0.06), dome-shaped, dorsally oval, and with anterior margin round and posterior margin somewhat attenuating from lateral view ("bicycle helmet" shaped); basal compound eyes anteroventrally pronounced, invisible to slightly visible from dorsal view, 0.52–0.66 mm in length, 0.29–0.36 mm in height, and 0.83–1.03 mm in distance between compound eyes (B/D = 3.11–4.91). **Thorax:** Forewings transparent, often with dark purplish brown markings at basal, central, and apical areas, 6.6–8.5 mm in length, and 2.2–3.2 mm in width; longitudinal veins light brown to dark purplish brown; crossveins light brown to dark purplish brown (basally and anteriorly located crossveins often darker and more heavily infuscated); crossveins C-Sc 12–19, Sc-R1 12–16, and R1-R2 10–14 in number; crossveins in stigmatic area not anastomosed; MP2 basally connected to MP1 and CuA; ICuI basally connected to CuA and CuP (angle between ICuI and crossvein [CuI-CuP] larger than angle between ICuI and crossvein [CuI-CuA]). Hindwings transparent, 1.3–1.9 mm in length, and 0.7–1.0 mm in width; longitudinal veins white to light yellow; crossveins C-Se, Sc-R1, and some in distal part sometimes dark brown; Sc ending relatively proximally (length of Sc / distance from base of Sc to apex of distal margin = 0.76); crossveins C-Sc 4–10, apically concentrated; costal margin round, with weakly developed costal projection at 1/3 basally to midlength, Rs 0.27–0.40 mm; R1 0.56–0.75 mm; MPs 0.20–0.31 mm; MP1 0.47–1.06 mm. Male forefemora 0.45–1.67 mm, foretibiae 2.22–2.83 mm, foretarsals segments 1, 2, 3, 4, and 5 0.10–0.13 mm, 0.68–0.97 mm, 0.70–0.93 mm, 0.55–0.70 mm, and 0.25–0.30 mm (foretarsal segment 2 > 3 > 4 > 5 > 1), and foreclaws 0.13–0.15 mm. Female foretibiae > forefemora > foretarsi. Claws dissimilar. **Abdomen:** Abdominal segment 9 with moderately to well developed posterolateral projections. Penes Y-shaped or V-shaped, relatively short or rudimentary (height of penis / height of forceps segment 1 = 0.08–0.31); forceps 3-segmented, with segment 1 0.63–0.68 mm, segment 2 0.06–0.09 mm, and segment 3 0.06–0.08 mm in length, forceps segment 1 basally broad and apically slender with relatively gradual constriction (constriction angle > 135–150°) at midlength to 2/3 apically; forceps segments 2 and 3 often indistinctly demarcated. Caudal filaments light yellow, often with purplish brown to dark brown stripes, often with maculation at joints; cerci 1.1–1.7x length of body; terminal filament as long as or slightly longer than cerci.

**Larva.** Male body length 6.2–9.8 mm; caudal filaments 5.5–10.0 mm. Female body length 6.8–11.2 mm; caudal filaments 6.3–13.2 mm. Body surface glassy and relatively less setose. General body color light yellow to light brown with dark purplish brown markings. **Head:** Head (Fig. 2) light brown, 1.27–1.40 mm in length and 1.59–1.75 mm in width. Male compound eyes dorsally 0.48–0.58 mm in width and 0.43–0.56 mm in distance between compound eyes (B/D = 0.74–1.10). Female compound eyes dorsally 0.20–0.24 mm in width and 1.10–1.35 mm in distance between compound eyes (B/D = 4.67–6.00). Antennae 4.0 mm in length, with whorls of hairlike setae at each segment. Cephalic tusks absent. Clypeus greatly developed, 0.38–0.45 mm in length, 0.68–0.90 mm in basal width, 0.64–0.83 mm in apical width, with pronounced sublateral tubercles, and with fields of 8–10 hairlike setae subapicolaterally and 4–15 hairlike setae subbasolaterally. Labrum slightly narrower than clypeus (0.92x width of clypeus), slightly wider distally (maximum length 0.28 mm; basal width 0.56 mm; distal maximum width 0.60 mm); dorsal surface moderately setose, with basal and subapical hairlike setal fringes (basal setae as long as subapical setae); subapical setal fringe laterally longer (setae curved forward) and without dense arrangement; anterior and lateral margins with dense hairlike setal row; anterior margin slightly concave, with prominent median tubercle, and with three pairs of rudimentary submedian denticles; ventral surface with dense hairlike setal field along anterior margin, with row of 15–18 stout setae on subanterior margins, and with fields of 10–15 hairlike setae at submedian areas. Mandible dorsolateral margins with very long hairlike setal row; ventral surface without setal row; inner incisors slightly smaller to as large as outer incisors; incisors with 2-

3 apical teeth and 0–4 lateral denticles; prostheca rudimentary, with well developed fringe. Hypopharynx superlinguae laterally curved and apically pointed. Maxillae with dense hairlike setal field, mixed with row of comblike setae almost entirely on galeolacinnial crown, without medioapical comblike seta, and with row of dense hairlike setae on medial margin; maxillary palp segment 1 0.26 mm, segment 2 0.29 mm, and segment 3 0.12 mm in length; segment 2 with sparse hairlike setae along outer margin; segment 3 indistinctly demarcated from segment 2, with moderately developed outer margin, and with moderately developed setal field along outer margin. Labial glossae dorsoventrally elongated and ventrally stalked, with dense hairlike setae; paraglossae with dense hairlike setal field dorsoapically; labial palp segment 1 0.38 mm, segment 2 0.27 mm, and segment 3 0.20 mm in length; segment 3 indistinctly demarcated from segment 2, apically pointed, with dense hairlike setal field along outer margin, and with stout setal row along inner margin. **Thorax:** Pronotum anterolateral margins round; lateral margins with row of sparse hairlike setae. Forefemora 1.15–1.50 mm, foretibiae 1.00–1.38 mm, foretarsi 0.43–0.55 mm, and foreclaws 0.15–0.25 mm in length (forefemora > foretibiae > foretarsi > foreclaws); forefemora with a few hairlike setae on dorsal and ventral surfaces, and with hairlike setal field along posterior margin (anterior margin with few hairlike setae); foretibiae with relatively sparse hairlike setae (filtering setae) dorsally and along outer margin (inner margin with few hairlike setae), with stout setal field (raking setae) 5/6 apically on inner margin; foretarsi with relatively sparse hairlike setae on dorsal and lateral surfaces; foreclaws apically darker, with rowed tiny teeth basally. Midlegs and hindlegs moderately setose; length femora > tibiae > tarsi > claws. **Abdomen:** Terga light yellow to light brown, mostly with broad submedian dark brown stripes, without long hairlike setal field along median line, and with hairlike setal row along lateral margins; abdominal segment 8–9 with moderately to well developed posterolateral projections. Sterna bare. Gills on abdominal segment 1–7, double; both lamellae with indistinct to distinct tracheae, with single apical filament; gill lamella inner part strongly expanded apicolaterally, and with fine setae on 1/10–1/2 apical margin; gill lamella outer part with fine setae on apically 1/10 to entire margin; apical filament attenuating, marginally with fine setae; gills 4 1.11–1.67 mm in length, 0.56–0.89 mm in width, 0.95–1.43 mm in filament length, with weakly to strongly developed apical expansion. Caudal filament segments with whorls of setae.

**Egg.** Egg (Fig. 5) oval; long axis 149 µm; short axis 100 µm. Color light yellow in nature, white in alcohol. Egg surface (Fig. 6) with ca. 116 knob-terminated coiled threads nearly evenly distributed throughout egg surface, with 0.4–2.1 µm tiny granules throughout egg surface; diameter of knob-terminated coiled threads 8.6–10.3 µm. Polar caps absent.

**Diagnosis.** The adults and larvae of *Ulmmerophlebia* can be distinguished from those of *Jappa* as specified above, in the diagnosis of *Jappa*. The larvae of *Ulmmerophlebia* can be distinguished from other leptophlebiid genera by the following combination of characters: possessing a prominent median tubercle anteriorly on the labrum, lacking cephalic tusks, and possessing a somewhat enlarged clypeal margin with sublateral tubercles and subapicolateral and subbasolateral hairlike setal fields.

**Species included.** *Ulmmerophlebia mjobergi* (Ulmer, 1916), *U. pipinna* Suter (1986), *Ulmmerophlebia* AV2 (Dean, 1999), *Ulmmerophlebia* AV3 (Dean, 1999), and *Ulmmerophlebia* AV5 (Dean, 1999).

**Distribution.** NSW, QLD, and VIC.

**Remarks.** *Deleatidium annulatum* Harker (1950) [Holotype: M; locality: NSW, Point Lookout, Serpentine; deposition: Australian Museum, Sydney (destroyed)] was considered as *Ulmmerophlebia* by Dean (1999), but the original description of the adult (e.g. wings) by Harker (1950) does not support placement in *Ulmmerophlebia* and the description of the larva (e.g. maxillary palp) by Harker (1954) does not meet the generic concept of *Ulmmerophlebia*.

*Atopopus spadix* Harker (1950) [Holotype: M; locality: NSW, Armidale; deposition: Australian Museum, Sydney (destroyed)] was also considered as

*Ulmerophlebia* by Dean (1999), but the original description of the adult (e.g. wings) by Harker (1950) does not support its placement in *Ulmerophlebia*.

#### DISCUSSION

In the phylogeny of the Atalophlebiinae (Leptophlebiidae), the genera *Jappa* and *Ulmerophlebia* have been hypothesized to constitute a basal clade, including *Hapsiphlebia*, *Atalophlebia*, *Atalomicrotia*, *Acanthophlebia*, *Aprionix*, and *Kalbaybaria* (Pescador and Peters, 1980; Towns and Peters, 1980, 1996; Campbell, 1993; Christidis, 2001). This clade, also known as the *Hapsiphlebia* lineage, was defined by the synapomorphies of lateral setae of the larval abdomen and incisor denticulation of the right mandible (Pescador and Peters, 1980).

From our comprehensive examinations of the species of *Jappa* and *Ulmerophlebia* and the species of related outgroup genera, we recognize an additional clade consisting of *Jappa* and *Ulmerophlebia*. This clade is defined by the synapomorphies of 1) a setose body, 2) gills marginally clothed with fine setae, 3) a median denticle anteriorly on the labrum, 4) submedian setal fields ventrally on the labrum, and 5) lateral spines on the larval abdomen.

*Jappa* and *Ulmerophlebia* are thus distinct monophyletic sister groups. *Jappa* is defined by the synapomorphies of 1) cephalic tusks, 2) a hairlike setal field ventrally on the mandible, 3) double rows of hairlike setae (filtering setae) dorsally on the tibiae, and 4) a long hairlike setal field along median abdomen. The cephalic tusks (Fig. 1) are unique in Leptophlebiidae. The cephalic tusks are arched and apically convergent and upward. They also bear basodorsal tubercles and setal fields at basomedial, subbasolateral, and apicodorsal areas. The body setation in *Jappa* is more specialized in having double rows of filtering setae on inner and outer margins of tibiae and broad setal fields on forefemora and along the median line of dorsal abdomen.

The monophyly of the *Ulmerophlebia* is supported by the synapomorphy of the unique "shovel-like" clypeal development (Fig. 2). The clypeus of *Ulmerophlebia* is flattened and somewhat elongated and possesses sublateral tubercles and hairlike setal fields in the subbasolateral and subapicolateral areas.

Although the adults of *Jappa-Ulmerophlebia* clade retain many plesiomorphic characters shared with other *Hapsiphlebia* lineage groups, the larvae are quite specialized as shown herein. The hairy body is associated with the fossorial habit of the members of *Jappa-Ulmerophlebia* clade and the cephalic tusks and shovel in *Jappa* and *Ulmerophlebia*, respectively, are evidently burrowing devices. In particular, the cephalic tusks of *Jappa* are analogous to the mandibular tusks of the Potamanthidae in Ephemeroidea, their Laurasian counterpart (Bae and McCafferty, 1991, 1995), in terms of functional morphology and burrowing behavior.

As evidenced above, the genera *Jappa* and *Ulmerophlebia* are distinct monophyletic groups that are here recognized at the generic level. Presumably, this will confirm with a strict phylogenetic classification if indeed the *Jappa-*

*Ulmerophlebia* clade, which is unique among Australian leptophlebiids being a burrowing mayfly group, is recognized as a distinct tribe as suggested for groupings under the subfamilies of Leptophlebiidae by Peters (1980). That classification can be adopted when the generic phylogenies of the Australian Leptophlebiidae are completed.

#### ACKNOWLEDGMENTS

We thank John Dean (Victoria EPA), Phil Suter (LaTrobe University), Richard Marchant (Museum of Victoria), Kimberi Pullan (ANIC, Canberra), Fred Govedich and Bonnie Bain (Monash University), Nigel Ainsworth (DPI, Victoria), and Dennis O'Dowd (Monash University) for their loans of material, field work assistance, and useful comments. We also thank anonymous reviewers for their useful comments and suggestions. This research was conducted during YJB's sabbatical leave to Monash University (September 2000–August 2001) that was supported by the Korea Research Foundation Grant (KRF-2001-013-G00007).

#### LITERATURE CITED

- Bae, Y. J. and W. P. McCafferty. 1991. Phylogenetic systematics of the Potamanthidae (Ephemeroptera). *Transactions of the American Entomological Society* 117:1-143.
- Bae, Y. J. and W. P. McCafferty. 1995. Ephemeroptera tusks and their evolution. pp. 377-405. *In*, L. Corkum and J. Ciborowski (Editors). *Current Directions in Research on Ephemeroptera*. Canadian Scholar's Publishing, Inc. Toronto, Canada. 478 pp.
- Campbell, I. C. 1990. The Australian mayfly fauna: composition, distribution and convergence. pp. 149-153. *In*, I. C. Campbell (Editor) *Mayflies and Stoneflies*. Kluwer Academic Publishers. Dordrecht, The Netherlands. 366 pp.
- Campbell, I. C. 1993. A new genus and species of leptophlebiid mayfly (Ephemeroptera: Leptophlebiidae: Atalophlebiinae) from tropical Australia. *Aquatic Insects* 15: 159-167.
- Christidis, F. 2001. Cladistic analysis of *Austrophlebiodes* and related genera (Leptophlebiidae: Atalophlebiinae). pp. 305-312. *In*, E. Dominguez (Editor). *Trends in Research in Ephemeroptera and Plecoptera*. Kluwer Academic / Plenum Publishers. New York, NY, U.S.A. 490 pp.
- Dean, J. C. 1999. Preliminary Keys for the Identification of Australia Mayfly Nymphs of the Family Leptophlebiidae. Cooperative Research Center for Freshwater Ecology. Identification Guide No. 20, Albury, New South Wales, Australia. 91 pp.
- Dean, J. C. and P. J. Suter. 1996. Mayfly Nymphs of Australia. A Guide to Genera. Cooperative Research Center for Freshwater Ecology, Identification Guide No. 7. Albury, New South Wales, Australia. 82 pp.
- Demoulin, G. 1955. Note sur deux nouveaux genres de Leptophlebiidae d'Australie (Ephemeroptera). *Bulletin et Annales de la Société Royale Entomologique de Belgique* 91:227-229.
- Eaton, A. E. 1871. A monograph on the Ephemeroidea. *Transactions of the Entomological Society of London* 1871:1-164.
- Eaton, A. E. 1884. A revisional monograph of recent Ephemeroidea or mayflies. Part II. *Transactions of the Linnean Society, Second Series, Zoology* 3:77-152.
- Edmunds, G. F., Jr., and W. P. McCafferty. 1996. New field observations on burrowing in Ephemeroptera from around the world. *Entomological News* 107:68-76.
- Harker, J. E. 1950. Australian Ephemeroptera. Part I. Taxonomy of New South Wales species and evaluation of taxonomic characters. *Proceedings of the Linnean Society of New South Wales* 71:1-34.

- Harker, J. E. 1954. The Ephemeroptera of Eastern Australia. Transactions of the Royal Entomological Society of London 105:241-268.
- Pescador, M. L. and W. L. Peters. 1980. Phylogenetic relationships and zoogeography of cool-adapted Leptophlebiidae (Ephemeroptera). pp. 43-56. In: J. F. Flannagan and K. E. Marshall (Editors) Advances in Ephemeroptera Biology. Plenum Press, New York, NY, U.S.A. 552 pp.
- Peters, W. L. 1980. Phylogeny of the Leptophlebiidae (Ephemeroptera), an introduction. pp. 33-41. In: J. F. Flannagan and K. E. Marshall (Editors) Advances in Ephemeroptera Biology. Plenum Press, New York, NY, U.S.A. 552 pp.
- Peters, W. L. and I. C. Campbell. 1991. Ephemeroptera (Mayflies). pp. 279-293. In: The Insects of Australia. Second Edition, Vol. 1. Melbourne Univ. Press, Carlton, Victoria, Australia. 542 pp.
- Riek, E. F. 1970. Ephemeroptera (Mayflies). pp. 224-240. In: The Insects of Australia. Melbourne University Press. Carlton, Victoria, Australia. 1029 pp.
- Skedros, D. G. and D. A. Polhemus. 1986. Two new species of *Jappa* from Australia (Ephemeroptera: Leptophlebiidae). The Pan-Pacific Entomologist 62:311-315.
- Suter, P. J. 1986. The Ephemeroptera (mayflies) of South Australia. Records of the South Australian Museum 19:339-397.
- Towns, D. R. and W. L. Peters. 1980. Phylogenetic relationships of the Leptophlebiidae of New Zealand (Ephemeroptera). pp. 57-69. In: J. F. Flannagan and K. E. Marshall (Editors). Advances in Ephemeroptera Biology. Plenum Press, New York, NY, U.S.A. 552 pp.
- Towns, D. R. and W. L. Peters. 1996. Leptophlebiidae (Insecta: Ephemeroptera). Fauna of New Zealand. 36:1-143.
- Ulmer, G. 1916. Results of Dr. E. Mjöberg's Swedish Scientific Expedition to Australia 1910-1913. 6. Ephemeroptera. Arkiv för Zoologi 10:1-18.
- Ulmer, G. 1920. Übersicht über die Gattungen der Ephemeropteren, nebst Bemerkungen über einzelne Arten. Stettiner Entomologische Zeitung 81:97-144.
- Williams, W. D. 1968. Australian Freshwater Life. The Invertebrates of Australian Inland Waters. Sun Books. Melbourne, Australia. 262 pp.

## A NEW SPECIES OF *PLATYCEPHALA* FROM CHINA (DIPTERA: CHLOROPIDAE: CHLOROPINAE)<sup>1</sup>

Shuwen An<sup>2</sup> and Ding Yang<sup>3</sup>

**ABSTRACT:** The genus *Platycephala* is newly recorded from Guangxi, with the description of a new species *P. guangxiensis*. Remarks on its relationships with the close species *P. zhejiangensis* Yang and Yang, 1995, are given.

**KEY WORDS:** Diptera, Chloropidae, Chloropininae, *Platycephala*, China.

The genus *Platycephala* Fallén belongs to the subfamily Chloropininae, and is characterized by the following features: Body large; head distinctly longer than high; frontal triangle occupying large part of frons, reaching anterior margin of frons with broadened apex; arista slender with short pubescence (Kanmiya, 1983). Until now the genus *Platycephala* contained seventeen species worldwide, of which eleven are known from the Oriental Realm (Sabrosky, 1977; Cherian, 1978; Kanmiya, 1983; Yang and Yang, 1994, 1995, 1997; An and Yang, 2003) and six from the Palearctic Realm (Nartshuk, 1984). Eight species are known from China (An & Yang, 2003). The major references dealing with *Platycephala* are as follows: Andersson (1977), Cherian (1978), Kanmiya (1983).

In this paper, one species of the genus *Platycephala* from Guangxi is described as new to science. Types are deposited in the Insect Collections of the China Agricultural University, Beijing.

### *Platycephala guangxiensis* An and Yang, NEW SPECIES (Figs. 1-8)

**Diagnosis:** Head triangular in profile, about 1.5 times as long as high. Frontal triangle trapezoidal, polished brownish yellow with two blackish lateral spots. Thorax black; propleuron brown with dark brown spot at posterior margin; pteropleuron blackish brown with pale upper part. Legs pale yellow.

**Description:** Male: Body length 5.5-6.5 mm, wing length 3.8-4.6 mm.

**Head** (Figs 1-2) blackish brown, triangular in profile, about 1.5 times as long as high; frons strongly produced beyond anterior level of eye, in profile about 0.35 times as long as long axis of eye; gena prominently broadened posterad, anteriorly strongly narrowed below eye; parafacial about 0.07 times as broad as long axis of eye. Frontal triangle trapezoidal, polished brownish yellow, occupying most of frons and reaching its anterior margin, semicircular anteriorly with two blackish lateral spots, and with many transverse grooves occupying median longitudinal area in front of ocelli; ocellar triangle black; area between eye and frontal triangle depressed and brownish. Occiput subshiny and punctured; gena and frons (in front of eye) brownish. Hairs and bristles on head brownish black. Antenna (Fig 3) brown with pale gray pollen; pedicel 0.8 times as long as flagellum; flagellum nearly rectangular, 1.6 times as long as wide, rounded apically, with blackish brown anterior and

<sup>1</sup> Submitted on May 15, 2003. Accepted on May 18, 2004.

<sup>2</sup> Department of Entomology, China Agricultural University, Beijing 100094, China. E-mail: shuwen\_an@hotmail.com.

<sup>3</sup> Department of Entomology, China Agricultural University, Beijing 100094, China. E-mail: dingyang@cau.edu.cn or dyangcau@yahoo.com.cn. Corresponding author.