## LISTS OF SPECIES

# Mayflies (Insecta: Ephemeroptera) of Yellowstone National Park, U.S.A.

## W. P. McCafferty

Department of Entomology, Purdue University, West Lafayette, Indiana, U.S.A. 47907. E-mail: mccaffer@purdue.edu

#### **Abstract**


The Ephemeroptera (Insecta) fauna of Yellowstone National Park consists of 46 species in 24 genera among eight families. These species are listed, and fifteen of the species (including collection data) are reported for the first time. Another 13 species have been taken adjacent to the park in Wyoming and Montana and noted as expected to occur in the park.

#### Introduction

Yellowstone National Park (Figure 1) was established as the world's first national park in 1872, and remains today the largest national park in the conterminous U.S.A. It comprises 898,714 ha, 91 % of which is in the northwestern corner of the state of Wyoming, with 7.6 % in adjacent Montana, and 1.4 % in adjacent Idaho. The roughly rectangular area is more than 100 km from north to south, and over 86 km from east to west. Approximately 10 % of the park is covered in water, including a variety of lakes and both high and low gradient streams and rivers. Although the climate has an average min/max temperature range of -13° C to -2° C in January, to 7° C to 27° C in July, some seasonal stream temperatures are moderated to various degrees because of geothermal water sources (Anonymous 1998).

Published reports of Ephemeroptera species from Yellowstone National Park have been scattered throughout the literature since the first report of a mayfly from there given by Eaton (1884) for what is now known as *Drunella grandis* (Eaton). Historically, 29 mayfly species have been reported from the park. Besides the Eaton report cited above, first reports for these species included eight in the 1920s, 11 in the 1930s, one in the 1940s, one in the 1950s, six in the 1960s, one in the 1980s, and one in the 2000s. This pattern is

typical of the explosion of Ephemeroptera data generated in North America in the 1920s and 1930s (primarily by James McDunnough and Jay Traver), which McCafferty (2001) has documented and referred to as the golden age of discovery of Ephemeroptera in North America. Yellowstone has been the type locality of four of these species: thermophilos (McDunnough) Acerpenna (McDunnough 1926), **Ecdyonurus** criddlei (McDunnough) (McDunnough 1927), Caudatella heterocaudata (McDunnough) (McDunnough 1929), and Rhithrogena futilis McDunnough (McDunnough 1934). Thus, with the exception of an informal record by Arbona (1980), and a record given by Slater and Kondratieff (2004), it has been nearly 40 years since significant new species records have been reported from the park. A few other species have been listed in various park lists or reports (grey literature). These are not treated here because identifications cannot be verified and because it is not the policy of Check List to include such species.



**Figure 1**. Yellowstone National Park; location within the U.S.A. and pertinent landmarks.

## LISTS OF SPECIES

New records for an additional 15 species of Ephemeroptera are presented below, based on the identification of materials donated to the Purdue Entomological Research Collection, Purdue University, West Lafayette, Indiana, U.S.A., by the late George Edmunds (University of Utah) in the 1990s. A checklist of all Yellowstone species is presented in alphabetical order by taxon. Newly reported species are asterisked within the list and

new record data are annotated for them. Previous species are annotated with a reference to their first, but not necessarily only, published Yellowstone record. All previous and new records are from the main portion of the park encompassed within the boundaries of Wyoming, including all of Park County and northern Teton County. A discussion of additional species that should eventually be found in the park follows the checklist.

## **Annotated checklist of species**

## Ameletidae

\*Ameletus bellulus Zloty, 1996: Larvae, Park County, no other locale data, 10-VII-1961, R. Muttkowski. *Ameletus sparsatus* McDunnough, 1931: Arbona (1980).

\*Ameletus validus McDunnough, 1923: Larvae, Park County, Antelope Creek, 2 miles south of Tower Junction, 17-VIII-1959, R. Muttkowski.

Ameletus velox Dodds, 1923: McDunnough (1934).

## **Baetidae**

\*Acentrella insignificans (McDunnough), 1926: Larvae, Teton County, Crawfish Creek, 10 miles south of West Thumb, 27-VI-1964, S. Jensen, J Richardson.

Acerpenna thermophilos (McDunnough), 1926: McDunnough (1926).

\*Baetis bicaudatus Dodds, 1923: Larvae, Park County, Blacktail Deer Creek, 7 miles east of Mammoth Hot Springs, 26-VI-1964, S. Jensen, J. Richardson.

\*Baetis flavistriga McDunnough, 1921: Larvae, Park County, Slough Creek, at Slough Creek Camp, 17-VIII-1959, G. Edmunds, W. Peters.

Baetis tricaudatus Dodds, 1923: Traver (1935).

Callibaetis ferrugineus (Walsh), 1862: Traver (1935) as C. fuscus Dodds, 1923.

\*Centroptilum conturbatum McDunnough, 1929: Larvae, Teton County, Crawfish Creek, 10 miles south of West Thumb, 27-VI-1964, S. Jensen, J. Richardson.

Diphetor hageni (Eaton), 1885: Traver (1935).

\*Fallceon quilleri (Dodds), 1923: Larvae, Teton County, Firehole River, 16 miles south of Madison Junction, 17-VIII-1959, G. Edmumds, W. Peters.

\*Heterocloeon anoka (Daggy), 1945: Larvae, Teton County, Firehole River, 6 miles south of Madison Junction, 16-VIII-1959, G. Edmunds, W. Peters.

## **Ephemerellidae**

Attenella margarita (Needham), 1927: Allen and Edmunds (1961a).

Caudatella heterocaudata (McDunnough), 1929: McDunnough (1929).

Caudatella hystix (Traver), 1934: Allen and Edmunds (1961b).

Drunella coloradensis (Dodds), 1923: Needham (1927).

Drunella doddsii (Needham), 1927: Needham (1927).

Drunella flavilinea (McDunnough), 1926: McDunnough (1935) as Ephemerella lapidula McDunnough, 1935.

Drunella grandis (Eaton), 1884: Eaton (1884).

Drunella spinifera (Needham), 1927: McDunnough (1934).

Ephemerella alleni Jensen and Edmunds, 1966: Jensen and Edmunds (1966).

Ephemerella aurivillii (Bengtsson), 1908: Vincent (1967).

## LISTS OF SPECIES

Ephemerella dorothea Needham, 1908: Allen and Edmunds (1965) as E. infrequens McDunnough, 1924.

Ephemerella excrucians Walsh, 1862: Needham (1927) as E. inermis Eaton, 1884.

Serratella tibialis (McDunnough), 1924: Traver (1934) as E. angusta Traver, 1934.

Timpanoga hecuba (Eaton), 1884: Allen and Edmunds (1959).

## **Ephemeridae**

Ephemera simulans Walker, 1853: Spieth (1940).

## Heptageniidae

Cinygmula mimus (Eaton), 1885: Slater and Kondratieff (2004).

Ecdyonurus criddlei (McDunnough), 1927: McDunnough (1927).

Epeorus albertae (McDunnough), 1924: McDunnnough (1929).

\*Epeorus deceptivus (McDunnough), 1924: Larvae, Park County, Tower Creek, at Tower Junction, 17-VIII-1959, G. Edmunds, W. Peters.

Epeorus grandis (McDunnough), 1924: Vincent (1967).

Epeorus longimanus (Eaton), 1885: Muttkowski and Smith (1929).

\*Heptagenia elegantula (Eaton), 1885: Larvae, Teton County, Crawfish Creek, 10 miles south of West Thumb, 27-VI-1964, S. Jensen, J. Richardson.

\*Heptagenia solitaria McDunnough, 1924: Larvae, Teton County, Crawfish Creek, 5-VI-1946, G Edmunds. Rhithrogena futilis McDunnough, 1934: McDunnough (1934).

Rhithrogena hageni Eaton 1885: Vincent (1967).

\*Rhithrogena morrisoni (Banks), 1924: Larvae, Park County, Soda Butte Creek, 1 mile east of Soda Butte, 17-VIII-1959, G Edmunds, W. Peters.

\**Rhithrogena robusta* Dodds, 1923: Larvae, Park County, Cub Creek, 15-VI-1949, G. Edmunds. *Rhithrogena undulata* (Banks), 1924: Traver (1935).

## Leptohyphidae

Tricorythodes explicatus (Eaton), 1892: McDunnough (1931).

\*Tricorythodes minutus Traver, 1935: Adults, Teton County, Madison River, 10-X-1947, F. Harmston.

## Leptophlebiidae

\*Leptophlebia cupida (Say), 1823: Adult, Teton County, Firehole River, nr Old Faithful, 21-VI-1949, G. Edmunds.

## **Siphlonuridae**

Siphlonurus occidentalis (Eaton), 1885: Traver (1935).

## **Discussion**

All species listed above are boreal northwestern or more widespread North American species (McCafferty 2006), and none show an unusual distribution within Yellowstone National Park. In addition to the 45 species listed above, there are another 13 species that I have seen in the Purdue Entomological Research Collections from just south of the park in Teton County, Wyoming, or in adjacent areas of Gallatin and Park Counties, Montana that may eventually be taken within the

These include Ameletus celer park proper. McDunnough, A. cooki McDunnough, majusculus Zloty, A. oregonensis McDunnough, similior McDunnough, vernalis A. McDunnough, Caenis youngi Roemhild, Centroptilum asperatum Traver, Cinygmula tarda (McDunnough), **Ecdyonurus** sinplicioides (McDunnough), Leptophlebia nebulosa (Walker), Parleptophlebia packii (Needham), and P. temporalis (McDunnough).

## LISTS OF SPECIES

## Acknowledgements

I thank R. P. Randolph (University of California at Davis) and A. V. Provonsha (Purdue University) for their assistance.

## Literature cited

- Allen, R. K. and G. F. Edmunds. 1959. A revision of the genus *Ephemerella* (Ephemeroptera: Ephemerellidae) I. The subgenus *Timpanoga*. Canadian Entomologist 91: 51-58.
- Allen, R. K. and G. F. Edmunds. 1961a. A revision of the genus *Ephemerella* (Ephemeroptera: Ephemerellidae). III. The subgenus *Attenuatella*. Journal of the Kansas Entomological Society 34: 161-173.
- Allen, R. K. and G. F. Edmunds. 1961b. A revision of the genus *Ephemerella* (Ephemeroptera: Ephemerellidae). II. The subgenus *Caudatella*. Annals of the Entomological Society of America 54: 603-612.
- Allen, R. K. and G. F. Edmunds. 1965. A revision of the genus *Ephemerella* (Ephemeroptera: Ephemerellidae). VIII. The subgenus *Ephemerella* in North America. Miscellaneous Publications of the Entomological Society of America 4: 244-282.
- Anonymous. 1998. Wyoming atlas & gazetteer. Yarmouth, DeLorme. 72 p.
- Arbona, F. L. 1980. Mayflies, the angler, and the trout. Tulsa, Winchester Press. 188 p.
- Eaton, A. E. 1883-88. A revisional monograph of recent Ephemeridae or mayflies. Transactions of the Linnaean Society of London, 2<sup>nd</sup> Series, Zoology 3: 1-352.
- Jensen, S. L. and G. F. Edmunds. 1966. A new species of *Ephemerella* from western North America (Ephemeroptera: Ephemerellidae). Journal of the Kansas Entomological Society 39: 576-579.
- McCafferty, W. P. 2001. The gentle quest: 200 years in search of North American mayflies. Pp.21-35. In E. Dominguez (ed.), Trends in Research in Ephemeroptera & Plecoptera. New York, Kluwer Academic/ Plenum Publishers.
- McCafferty, W. P. 2006. Mayfly central: Mayflies of North America. Electronic database accessible at http://www.entm.purdue.edu/entomology/research/mayfly/mayfly.html.

- Purdue University, West Lafayette, Indiana, U.S.A. Captured on 31 October 2006.
- McDunnough, J. 1926. New Canadian Ephemeridae with notes. IV. Canadian Entomologist 58: 296-303.
- McDunnough, J. 1927. A new *Heptagenia* from the Yellowstone region (Ephemeroptera). Canadian Entomologist 59: 261.
- McDunnough, J. 1929. Notes on North American Ephemeroptera with descriptions of new species, II. Canadian Entomologist 61: 169-180
- McDunnough, J. 1931. New North American Caeninae with notes (Ephemeroptera). Canadian Entomologist 63: 254-268.
- McDunnough, J. 1934. New species of North American Ephemeroptera. IV. Canadian Entomologist 66: 154-164, 181-188.
- McDunnough, J. 1935. Notes on western species of Ephemeroptera. Canadian Entomologist 67: 95-104.
- Muttkowski, R. A. and G. M. Smith. 1929. The food of trout stream insects in Yellowstone National Park. Roosevelt Wildlife Annals of the Roosevelt Forestry Experiment Station of the New York State College of Forestry at Syracuse University 2: 241-263.
- Needham, J. G. 1927. The Rocky Mountain species of the mayfly genus *Ephemerella*. Annals of the Entomological Society of America 20: 107-117.
- Slater, J. and B. C. Kondratieff. 2004. A review of the mayfly genus *Cinygmula* McDunnough (Ephemeroptera: Heptageniidae) in Colorado. Journal of the Kansas Entomological Society 77: 121-126.
- Spieth, H. T. 1940. Studies on the biology of the Ephemeroptera. II. The nuptial flight. Journal of the New York Entomological Society 48: 379-390.
- Traver, J. R. 1934. New North American mayflies (Ephemerida). Journal of the Elisha Mitchell Scientific Society 50: 189-254.
- Traver, J. R. 1935. Part II, Systematic. Pp 237-739. In J.G. Needham, J.R. Traver, and Y.C Hsu (eds.), The biology of mayflies with a systematic account of North American species. Ithaca, Comstock Publishing.
- Vincent, E. R. 1967. A comparison of riffle insect populations in the Gibbon River above and below the Geyser Basins, Yellowstone

## LISTS OF SPECIES

National Park. Limnology and Oceonography 12: 18-26.

Received November 2006 Accepted December 2007 Published online January 2007