



Ephemeroptera and Plecoptera biodiversity in central Patagonia, Chubut province, Argentina

PABLO PESSACQ¹ & MARÍA LAURA MISERENDINO²

Laboratorio de Investigación en Ecología y Sistemática Animal (LIESA), Universidad Nacional de la Patagonia San Juan Bosco, Sarmiento 849, 9200. Esquel, Chubut, Argentina. E-mail: ¹pablopessacq@yahoo.com.ar; ²lmiserendino@liesa.org.ar

Abstract

Aquatic insects in Central Patagonia (Andean region), Argentina, have received poor and fragmentary attention. In this paper we present the first preliminary inventory of Plecoptera and Ephemeroptera taxa for the area and include seasonal records of adult stages. We incorporate a set of environmental features for the sampling sites and biotopes as well as the main characteristics of the phytogeographical and biogeographical areas in which the species were found. Twelve species and eight genera of mayflies and eleven species and five genera of stoneflies are recorded for the first time in Chubut province. The stoneflies *Neofulla biloba* (Aubert), *Alfonsoperla flinti* McLellan & Zwick, *Austronemoura eudoxiae* Froehlich, *Chilenoperla semitincta* Illies, *Neonemura illiesi* Zwick and *Potamoperla testacea* Vera, and the mayflies *Andesiops ardua* (Lugo-Ortiz & McCafferty), *Chaquihua bullocki* Navás. And the family Oniscigastridae are new records for Argentina. The genera *Americabaetis* Kluge, *Camelobaetidius* Demoulin and *Traverella* Edmunds are confirmed for first time in the Andean region (Central Patagonia province) and four Plecoptera and two Ephemeroptera species constitute new records for the Patagonian steppe. We also include two new records for Ephemeroptera and Plecoptera for the Santa Cruz province. We reinforce the idea that patagonian aquatic environments have a highly specialized and complex insect fauna which can be correlated with biogeographical aspects but also with a great environmental heterogeneity.

Key words: new records, inventory, Patagonia, Ephemeroptera, Plecoptera

Introduction

Knowledge of Ephemeroptera and Plecoptera fauna in Argentina is still poor and fragmentary (Domínguez 1998, Muzón & Bachmann 1998). This is certainly true for the Central and the Subandean Patagonia (Patagonian subregion) in Chubut province, these areas had received little sampling effort. As a result of different research projects of the Laboratorio de Investigación en Ecología y Sistemática Animal (LIESA) several lotic environments in the area have been the object of intense and systematic collections (Miserendino 1996, 2001, Miserendino & Pizzolón 2000, 2004, Hollmann & Miserendino 2006). Even so, wide areas in the southern region, the Valdivian forest, and most of the steppe remain almost unexplored.

At present, 25 mayfly species within 13 genera have been recorded for the Argentinean Subantarctic and Patagonian subregions (Andean region) (Domínguez *et al.* 1994, Muzón & Pessacq 2001, Nieto 2004). Sixteen of those species (in 8 genera) have been recorded in the Chubut province (Domínguez *et al.* 1994, Miserendino & Pizzolón 2000, Miserendino 2001, Nieto 2004).

Stoneflies are represented by 36 species within 27 genera (McLellan *et al.* 1990, 2006, Muzón & Bachmann 1998, Nelson 1973), of which 17 (within 15 genera) have been recorded in the Chubut province (Aubert 1960, Illies 1963, McLellan *et al.* 2006, Miserendino 1996, Miserendino & Pizzolón 2000).

Material and methods

The present contribution is the result of several surveys carried out in 2005, 2006 and 2007 at 41 sampling sites (Table 1). The study area included the Middle West of Chubut province, the Chubut River basin, and the Santa Cruz River in Santa Cruz province, located within the Patagonian and Subantarctic subregions of the Andean region (Morrone 2006) (Fig. 1).




FIGURE 1. Sampling sites in Chubut province. Code of sampling sites in Table 1. P.N.: Los Alerces National Park.

For larval collections Surber, kick-net, and D-frame nets were used, whereas for adults aerial nets, pitfall traps, light traps and Malaise traps were employed (Hauer & Resh 1996).

An environmental characterization for most of the sampling sites was performed. For that purpose, we compiled information previously published (Miserendino 2001, 2006, Miserendino & Pizzolón 2004) and new data. Substrate composition was estimated as percentage of boulder, cobble, gravel, pebble, and sand using a 1-m² grid (Gordon *et al.* 1994). Stream order was obtained from Coronato & Del Valle (1988). Surface current speed was obtained by timing a bobber (average of 3 times) as it moved over a distance of 10 meters (Gordon *et al.* 1994). At each site we measured wet width and bank full width.

The phytogeographical and biogeographical description was based on León *et al.* (1998) and Morrone (2006) respectively. The species composition of the riparian corridor was analyzed and main native and exotic species recorded.

Results

Environmental Characterization: The main environmental features of the sampling sites are presented in Table 2. This gives a picture of the diversity of watercourses visited during the study. Elevation of sampling sites ranged between 3 and 820 m. Stream order was between 2 and 6. Conductivity values ranged between 9 and 306 $\mu\text{S}\cdot\text{cm}^{-1}$. As expected, mountain streams showed lower conductivity values than those in the steppe. Sub-

strate size was similar across most mountain rivers, being mainly comprised of boulders and cobbles. Substrate size in steppe streams was finer, consisting mostly of cobbles, gravel, and sand.

TABLE 1. Sampling locations for Chubut and Santa Cruz provinces, Argentina. Sites 1-6, 8-19 and 26-30 were sampled on several occasions between May 2005 and March 2006, sites 20-25 were visited between October 2006 and April 2007. RN: Ruta Nacional, RP: Ruta Provincial.

Chubut Province						
Site	Water Body	Locality	Other descriptor	Lat.	Long.	Dates
1	Golondrinas Str.	El Hoyo	Intersection RP 258	41°59'33"	71°33'25"	
2	Nameless Str.	Puerto Patriada	trib. Epuyen L.	42°08'17"	71°31'56"	
3	Las Minas Str.	Cholila	Pre-urban reach	42°31'03"	71°25'13"	
4	Las Minas Str.	Cholila	Urban reach	42°30'32"	71°25'34"	
5	Esquel Str.	Nr. Esquel	Road access to "La Hoya"	42°50'28"	71°15'55"	
6	Esquel Str.	Nr. Esquel	Pre-urban reach	42°53'48"	71°16'03"	
7	Percey R.	Alto Río Percey	15 km from Esquel	45°51'17"	71°25'23"	4-26/III/07
8	Percey R.	Nr. Trevelin	2 km upstream Trevelin	43°04'22"	71°28'14"	
9	Blanco Str.	Nr. Los Cipreses	At RN 259 intersection	43°10'38"	71°36'30"	
10	Baggilt Str.	Nr. Los Cipreses	At RN 259 intersection			
11	Nameless Str.	Nr. Trevelin	RP 17, Percey R. trib.	43°12'02"	71°24'01"	
12	Nant y Fall Str.	Nr. Trevelin	At RP 17 intersection	41°13'24"	71°25'17"	
13	Nameless Str.	Nr. Los Ñires farm	RP 17, Río Frío trib.	43°1'33"	71°26'36"	
14	Río Frío	Los Ñires farm	Farm on SR 17	43°29'56"	71°20'22"	
15	Nameless Str.	Nr. los Ñires farm	RP 17, Río Frío trib.	43°21'43"	71°30'22"	
16	Glyn Str.	Nr. Cerro Centinela	Nearby RP 17	43°27'34"	71°33'25"	
17	Carbón Str.	Nr. Corcovado	Pre-urban reach	43°32'11"	71°28'21"	
18	Cabeza de Vaca Str.	Nr. Corcovado	At RP 17 intersection	43°29'58"	71°20'51"	
19	Carbón Str.	Corcovado	Urban reach	43°32'07"	71°28'11"	
20	Corcovado R.	Nr. Corcovado	El Palenque farm, RP 44			
21	Poncho Moro Str.	Nr. Poncho Moro farm	RP 44 intersection			
22	El Comisario Str.		RP 44 intersection	43°44'45"	71°23'32"	
23	Nameless Str.		RP 44 & Corcovado trib.	43°44'36"	71°23'35"	
24	Corcovado R.		Outlet of Vintter L.	43°53'54"	71°25'28"	27/IV/07
25	Chubut R.	Fofó Cahuel	At RP 4	42°26'07"	70°30'35"	
26	Chubut R.	Piedra Parada	At RP 12	42°40'25"	70°05'1"	
27	Chubut R.	Paso Berwin	At RN 25	43°37'15"	68°57'05"	
28	Chubut R.	Las Plumas	At RN 25 intersection	43°43'12"	67°17'21"	
29	Chubut R.	Valle Los Altares	At RN 25	43°53'29"	68°23'5"	
30	Chubut R.	28 de Julio	Dwnstr. Ameghino dam	43°22'01"	65°53'17"	
31	Coronado Str.	Los Alerces Nat'l Pk.	West side Rivadavia L.	42°37'60"	71°41'55"	12/III/07
32	Colegual Str.	Los Alerces Nat'l Pk.	Confluence Rivadavia R.	42°40'27"	71°41'55"	26/XI/06
33	Nameless Str.	Los Alerces Nat'l Pk.	N of Centinela str, RN 71	42°29'06"	71°23'21"	29/III/07
34	Nameless Str.	Los Alerces Nat'l Pk.	Pucon Pai camping., RN 71	42°29'44"	71°23'21"	29/III/07

35	Irigoyen Str.	Los Alerces Nat'l Pk.	Irigoyen Fall, RN 71	42°30'57"	71°21'43"	29/III/07
36	Nilson R.	Los tres leones farm	2 km from Pico R.	44°15'06"	71°40'37"	29/I/07
37	Nameless str.	Nr. Río Pico	Access road to L. 2			28/IV/07
Santa Cruz Province						
38	Santa Cruz R.	Charles Für	At RP 9	50°15'35"	71°53'40"	V/05
39	Santa Cruz R.	Condor Cliff	At RP 9	50°13'07"	70°55'52"	V/05
40	Santa Cruz R.	Barrancosa	At RP 9	50°30'30"	70°10'41"	V/05
41	Santa Cruz R.	Cte Luis Piedrabuena	At RP 63	50°0'48"	69°0'32"	V/05

TABLE 2. Environmental features of the sampling sites. Phytogeographical units codes: 1) Subantarctic province. 2) Patagonic province: Subandean district: a) Herbaceous steppe, occidental district: b) Herbaceous/shrub like steppe, central district c) Barren steppe d) Shrub steppe of *Chuquiraga Avellanadae* e) Shrub steppe of *Junellia tridentis*. Ach: *Austrocedrus chilensis*, Nd: *Nothofagus dombeyi*, Na: *N. antarctica*, Np: *N. pumilio*, Mb: *Maytenus boaria*, S: *Salix spp.*, Pn: *Populus nigra*, Pa: *P. alba*, Pm: *Pseudotsuga mentziesii*, Pc: *Pinus contorta*. Boul: boulder, Co: cobble, Pe: pebble, Gra: gravel, sand.

Site	Stream order	Bank Width (m)	Substrate	Current m/sec	Conductivity uS/sec	Elevation m	Phytographic Units	Riparian Veg.
1	2	4.2	Co/Sand	2.6	126	361	1	Pm, Pc
2	1	1.8	Gra/Sand	3	76	495	1	Pm
3	2	3.25	Co/Boul	0.93	164	573	1	S
4	2	9	Pe/Co	3	72	555	1	S
5	3	10.8	Boul/Co	3	105	721	2a	Mb, Na
6	4	9	Co/Pe	0.9	184	610	2a	S, Na
7	5	80	Boul/Co	1	80	600	1	S
8	5	100	Boul/Co	1.11	83	360	2a	S
9	2	35	Boul/Co	0.4	16.8	450	1	Na
10	3	20	Boul/Co	0.6	21.6	460	1	Np, Sal
11	2	0.9	Gra/Sand	0.1	84	730	1	Na
12	3	31.5	Gra/Sand	1.45	113	690	1	Na
13	2	3.5	Co/Pe	3	9	670	1	Na
14	4	22.5	Sand/Gra	0.9	98	670	1	-
15	3	14.2	Boul/Co	1.5	87	700	1	-
16	2	5.66	Co/Pe	0.1	69	615	1	Na
17	3	10.5	Boul/Co	1.73	93	380	1	Pn, Pa
18	2	6.88	Co/Gra	1.06	63	700	1	Na, S
19	3	23.5	Boul/Co	1.35	84	400	1	S
20	4	50	Boul/co	1	-	430	1	S
21	3	12	Boul/Co	1	23.7		1	Na, Ach
22	2	4	Boul/Co	1.03	26	820	1	Na, Np
23	3	9	Boul/Co	0.8	29	770	1	Na, Np
24	4	30	Boul/Co	1.9	-	928	1	Na, Np
25	4	80	Pe/Gra	1.1-2.1	99	500	2a	S
26	5	200	Co/Pe	0.9-1.2	148.7	440	2b	S
27	5	160	Co/Sand	0.5-1.6	185.2	308	2d	S

28	6	200	Pe/Gra/Sand	0.8-1.2	187.2	243	2d	S
29	6	300	Boul/Co	1.7-1.9	189.5	300	2c	S
30	6	70	Co/Pe	0.7-0.9	196	70	2c	S, Pn, Pa
31	3	15	Co/Gra	0.8	-	541	1	Nd, Ach, S
32	3	25	Boul/Co	1.6	26.1	527	1	Nd, Ach
33	3	25	Boul/Co	0.7	41.2	580	1	Nd, Ach
34	3	3	Boul/Co	1.1	22.3	570	1	Nd, Ach, Ch
35	3	7	Boul/Co	1.3	26.1	570	1	Nd, Ach, Ch
36	3	10	Boul/Co	1.4	-	503	1	Nd, S
37	3	7	Gra/Co	0.2	-	-	1	Nd
38	6	150	Co/Pe	0.35	-	170	2e	-
39	6	200	Boul/Co	1.2	-	120	2e	-
40	6	230	Pe/Gra	0.5	-	82	2e	-
41	6	140	Pe/Gra	1.43	-	3	2e	-

Ephemeroptera and Plecoptera analysis: This contribution increases to 24 the number of mayfly species that are known for the Chubut Province (within 15 genera), including the first records for Argentina of *Andesiops ardua* (Lugo-Ortiz & McCafferty 1999), *Chaquihua* Demoulin, 1955b and the family Onis-cigastridae. Moreover, the genera *Camelobaetidius* Demoulin, 1966, *Americabaetis* Kluge, 1992 and *Traver-ella* Edmunds, 1948 are recorded for the first time from the entire Andean Region (Table 3).

TABLE 3. Ephemeroptera and Plecoptera species recorded for Chubut province (including data from Santa Cruz River, Santa Cruz province). Site of collection, comments, previous records, and main literature are indicated when the taxa represent a new record. CH: Chubut province, NQ: Neuquén province, RN: Río Negro province, SC: Santa Cruz province. Records of *Chaquihua* & *Siphonella* are in agreement with Domínguez *et al.* (2001) who mention the genus for southern Argentina in their generic key without further information.

	Sites	Comments	Previous records	References
EPHEMEROPTERA				
Ameletopsidae				
<i>Chaquihua bullocki</i> (Navás 1930)	1, 2, 16	New generic record, Argentina	Subant. subreg., S Chile	Mercado & Elliot 2004
<i>Chiloporter eatoni</i> Lestage 1931				Domínguez 1998
Baetidae				
<i>Americabaetis alphus</i> Lugo-Ortiz & McCafferty 1996	25, 26, 29	New generic record, Andean reg.	S Am. trans. zone, Neotrop. reg., Argentina	Lugo-Ortiz & Mc Cafferty 1996
<i>Andesiops ardua</i> (Lugo-Ortiz & McCafferty 1999)	29	New record, Argentina	Subant. subreg., S Chile	Lugo-Ortiz & Mc Cafferty 1999, Nieto 2004
<i>A. torrens</i> (Lugo-Ortiz & McCafferty 1999)				Nieto 2004
<i>Camelobaetidius penai</i> (Traver & Edmunds 1968)	25, 26, 29	New generic record, Andean reg.	S Am. trans. zone reg., Neotrop. reg., Argentina	Domínguez 1998
Caenidae				
<i>Caenis</i> sp.	3, 4, 5, 7, 25, 26, 28	New family record, Andean reg.	Neotrop. reg.	Domínguez <i>et al</i> 1994, Miserendino & Pizzolón 2000
Leptophlebiidae				

<i>Archethraulodes spatulus</i> Pesca- dor & Peters 1982	21	New generic record, Chubut prov.	Subant.& Patag. subreg., S Chile & Argentina (NQ)	Domínguez 1998
<i>Dactylophlebia carnulenta</i> Pescador & Peters 1980	18, 20-22, 37	New generic record, Chubut prov.	Subant. & Patag. sub- regs., S Chile & Argen- tina (NQ)	Domínguez 1998
<i>Meridialaris chiloeensis</i> (Demoulin 1955a)	38, 40, 41	New record, Patago- nia steppe & Santa Cruz prov.	Subant. & Patag. sub- reg.s, S Chile & Argen- tina (CH, NQ, RN)	Domínguez 1998
<i>M. diguillina</i> (Demoulin 1955c)				Domínguez 1998
<i>M. laminata</i> (Ulmer 1920)	5, 6, 7, 8, 11,12,13,1 4	1st confirmed record, Chubut prov	Subant.& Patag. subreg., S Chile & Argentina (NQ, RN)	Domínguez 1998, Miserendino & Pizzolón 2000
<i>Nousia bella</i> Pescador & Peters 1985	9, 10, 12, 20, 21, 37	1st confirmed record, Chubut prov	Subant. & Patag. subreg., S Chile & Argentina (NQ, RN)	Domínguez 1998, Miserendino 2001
<i>N. crena</i> Pescador & Peters 1985				Domínguez 1998
<i>N. delicata</i> Navás 1918b				Domínguez 1998
<i>N. minor</i> (Demoulin 1955c)				Domínguez 1998
<i>Penaphlebia vinosa</i> (Demoulin 1955c)	1	1st confirmed record, Chubut prov.	Patag. subreg., Argentina (NQ, RN)	Domínguez 1998, Miserendino 2001
<i>P. chilensis</i> (Eaton 1883)	26	New record, Patago- nia steppe	Subant. & Patag. sub- regs., S Chile & Argen- tina (CH, NQ, RN)	Domínguez et al 1994
<i>P. flavidula</i> Pescador & Peters 1991				Domínguez 1998
<i>P. fulvipes</i> (Needham & Murphy 1924)				Domínguez 1998
<i>Rhigotopus andinensis</i> Pescador & Peters 1982				Domínguez 1998
<i>Traverella</i> sp.	25-30	New generic record, Andean reg.	S Am. trans. zone reg., Neotrop. reg., Argentina	Domínguez 1998
Nesameletidae				
<i>Metamonius anceps</i> (Eaton 1885)				Domínguez 1998
Oniscigastridae				
<i>Siphonella</i> sp.	7, 15, 16, 32	New family record, Argentina	Subant. subreg., S Chile	Needham & Murphy 1924, Pictet 1894
PLECOPTERA				
Eustheniidae				
<i>Neuroperla schedingi</i> (Navás 1929)				Illies 1960b, Miserendino 1996
Diamphipnoidae				
<i>Diamphipnopsis samali</i> Illies 1960b	1, 2	New generic record, Chubut prov.	Subant. & Patag. sub- regs., S Chile & Argen- tina (NQ, RN)	Illies 1960b, Wais et al 1987
Austroperlidae				
<i>Klapopteryx kuscheli</i> Illies 1960a	38-41	New record, Patago- nia steppe & Santa Cruz prov.	Subant. & Patag. sub- regs., S Chile & Argen- tina (CH, NQ, RN)	Illies 1960a, McLellan 2001

Table 3. Continued

	Sites	Comments	Previous records	References
<i>Penturoperla barbata</i> Illies 1960a				Illies 1960a, Miserendino 1996
Griopterygidae				
<i>Alfonsoperla flinti</i> McLellan & Zwick 2007	15, 22,23	New generic record, Argentina	Subant. subreg., S Chile	McLellan & Zwick 2007
<i>Antarctoperla michaelsoni</i> (Klapálec 1904)	28, 30, 38-41	New record, Patagonia steppe	Subant. & Patag. subregs., S Chile & Argentina (CH, NQ, RN, SC)	Illies 1963, Miserendino 2001
<i>Aubertoperla illiesi</i> (Froehlich 1960)				Illies 1963
<i>Araucanioperla bullocki</i> (Navás 1933)				Illies 1963
<i>Ceratoperla fazi</i> (Navás 1934)	22, 23	New record, Chubut prov	Subant. & Patag. subregs., S Chile & Argentina (RN)	Illies 1963
<i>Chilenoperla semitincta</i> Illies 1963	5	1st confirmed record, Argentina	Subant. subreg., S Chile	Illies 1963, Miserendino 2001, Miserendino & Pizzolón 2000
<i>Limnoperla jaffueli</i> (Navás 1928)	29, 38-41	New record, Patagonia steppe	Subant. & Patag. subregs., S Chile & Argentina (CH, NQ, RN)	Illies 1963, Miserendino 2001
<i>Notoperla archiplatae</i> (Illies 1958)				Illies 1958
<i>N. fasciata</i> McLellan 2006				McLellan 2006
<i>N. magnaspina</i> McLellan 2006				McLellan 2006
<i>Notoperlopsis femina</i> Illies 1963	26	New record, Patagonia steppe	Subant. & Patag. subregs., S Chile & Argentina (CH)	Illies 1963, McLellan et al 2006
<i>Pelurgoperla personata</i> Illies 1963				Illies 1963
<i>Potamoperla myrmidon</i> (Mabille 1891)				Illies 1963
<i>P. testacea</i> Vera 2006	22, 23	New record, Argentina	Subant. subreg., S Chile	Vera 2006
<i>Rhithroperla rossi</i> (Froehlich 1960)				Froehlich 1960, Illies 1963
<i>Senzilloides panguipullii</i> (Navás 1928)				Illies 1963
Notonemouridae				
<i>Austronemoura chilena</i> Aubert 1960	15, 21 22, 33-35	New generic record, Chubut prov	Subant. subreg., S Chile & Patag. subreg., Argentina (RN)	Aubert 1960, Illies 1961
<i>A. eudoxiae</i> Froehlich 1960	15	New record, Argentina	Subant. subreg., S Chile	Froehlich 1960, Illies 1961
<i>Neofulla biloba</i> (Aubert 1960)	35	New generic record, Argentina	Subant. subreg., S Chile	Aubert 1960, Illies 1961
<i>Neonemura barrosi</i> Navás 1919	21, 31	New generic record, Chubut prov	Subant. subreg., S Chile & Patag. subreg., Argentina (RN, NQ)	Illies 1961, Wais et al 1987
<i>N. illiesi</i> Zwick 1972	10	New record, Argentina	Subant. subreg., S Chile	Zwick 1972

Udamocercia frantzi Illies 1961 5, 21, 22, New generic record, Subant. subreg., S Chile Illies 1961, Wais et al 1987
24, 37 Chubut prov & Patag. subreg., Argentina (NQ, RN)

Perlidae

Kempnyella genualis (Navás 1918b) Illies 1964

Pictetoperla gayi (Pictet 1841) Illies 1964

The number of stonefly species now known from Chubut Province stands at 28 (in 23 genera). Moreover, The genus *Neofulla* Claassen, 1936, *Alfonsoperla flinti* McLellan & Zwick, 2007, *Austronemoura eudoxiae* Froehlich, 1960, *Neonemura illiesi* Zwick, 1972, *Chilenoperla semitincta* Illies, 1963 and *Potamperla testacea* Vera, 2006 are recorded for the first time in Argentina.

Additionally, four species of stoneflies and two species of mayflies are recorded for the first time in Patagonia steppe. For Santa Cruz Province a new Ephemeroptera specific record and a new Plecoptera generic record are included. Table 3 provides a summary of the Ephemeroptera and Plecoptera inventory for Chubut province, including site of collection, comments, previous records and main references for every new record.

Discussion

The emergence periods of Ephemeroptera and Plecoptera species are predictable and well known in the northern hemisphere and in some southern hemisphere regions (Hynes & Hynes 1975, Scrimgeour 1991, Richardson 2001). Unfortunately, literature about emergence periods of mayflies and stoneflies in Patagonia is limited to that of Miserendino & Hollmann (2005) and Hollmann & Miserendino (2008). The emergence data presented in this paper are not the result of a systematic sampling, nevertheless, the capture dates of adults constitute new information for the species in the area (Table 4). Mayflies emerged nearly exclusively in southern hemisphere in spring and summer, with the minor exception of *Nousia delicata* Navás, 1918a, which was also present in early autumn. *Meridialaris chiloeensis* (Demoulin 1955a) was the first mayfly to emerge, beginning in early October, which is consistent with reports by Hollmann & Miserendino (2006) who observed a bivoltine pattern with spring and summer generations for this species.

Some stoneflies began emergence in late winter, but the majority waited until spring, with several emerging throughout the summer months. Adults of the three Notonemouridae genera *Austronemoura* Aubert, 1960, *Neofulla* and *Udamocercia* Enderlein, 1909 were found mainly by the end of summer and beginning of autumn. Most Gripopterygidae emerged during spring and summer months. *Notoperlopsis femina* Illies, 1963 appeared only in October, while *Ceratoperla fazi* (Navás 1934) and *Pelurgoperla personata* Illies, 1963 appeared to be early autumnal species.

The most abundant and ubiquitous Ephemeroptera species were *Meridialaris chiloeensis*, *M. diguillina* (Demoulin 1955c) and *Andesiops torrens* (Lugo-Ortiz & McCafferty 1999), whereas *Antarctoperla michaelsoni* (Klapalek 1904), *Aubertoperla illiesi* (Froehlich 1960), *Limnoperla jaffueli* (Navás 1928) and *Klapopteryx kuscheli* were the most frequently collected Plecoptera species. These qualitative abundances and generalized distributions for both mayflies and stoneflies are in agreement with previous reports in the area (Miserendino & Pizzolón 2000).

The records of *Camelobaetidius*, *Americabaetis* and *Traverella* were unexpected. Their southern distributional limit was the South American transition zone and the Neotropical region, in the Argentinean provinces of Córdoba and Tucumán, about 1000 km north of our records. These records, together with that of *Leptohyphes* Eaton, 1882 (Ephemeroptera), in Somuncurá Plateau, Río Negro province (Muzón & Pessacq 2001) suggest a disjunct distributional pattern with interesting biogeographical issues. However, these patterns must be corroborated with extensive sampling in intermediate areas.

TABLE 4. Months and seasons of collection for those species where adults were observed. *Indicates a low quantity of specimens collected at only one sampling site and date. (x) Indicates only one or two specimens collected.

Season Month	Winter		Spring			Summer			Autumn			
	Jul.	Ag.	Sept.	Oct.	Nov.	Dec.	Jan.	Feb.	Mar.	Apr.	May	Jun.
Ephemeroptera												
<i>Penaphlebia chilensis</i>				x								
<i>Meridialaris chiloensis</i>				x	x			x	x			
<i>Nousia delicata</i>				x	x		x	x			x	
<i>Andesiops ardua</i>								x				
<i>Archethraulodes</i> sp. *								x				
<i>Meridialaris laminata</i>					x							
<i>Meridialaris diguillina</i>								x	x			
<i>Metamonius anceps</i> *									x			
<i>Siphonella</i> sp.									x			
Plecoptera												
<i>Austronemoura chilena</i>										x		
<i>Antarctoperla michaelsoni</i>			x	x	x	x	x	x	x			
<i>Aubertoperla illiesi</i>			(x)	x	x	x	x	x	x			x
<i>Neonemura barrosi</i>			(x)					(x)	x			
<i>Notoperlopsis femina</i>				x								
<i>Klapopteryx kuscheli</i>				x	x	x				x		
<i>Limnoperla jaffueli</i>				x	x	x	x	x	x			
<i>Senzilloides panguipulli</i>				x						x		
<i>Potamoperla testacea</i>					x	x						
<i>Neofulla biloba</i> *										x		
<i>Pelurgoperla personata</i>										x		
<i>Penturoperla barbata</i> *										x		
<i>Rhithroperla rossi</i> *										x		
<i>Udamocercia frantzi</i>										x	x	
<i>Austronemoura eudoxiae</i> *										x		
<i>Ceratoperla fazi</i>										x	x	

Some taxa were rare or occasionally recorded, as was the case for *Austronemoura eudoxiae*, *Neofulla biloba* (Aubert 1960), *Neonemura illiesi*, *Potamoperla testacea*, *Alfonsoperla flinti* (Plecoptera) and *Archethraulodes spatulus* Pescador & Peters, 1982 (Ephemeroptera). These taxa were represented by only a few specimens at one sampling site. The gripopterygid *Potamoperla myrmidon* (Mabille 1891) was the only species that appeared through all sampling sites in Chubut River (Miserendino 2006). Similarly, the austroperlid *Klapopteryx kuscheli* was found in all the sampling sites in Santa Cruz River (province of Santa Cruz) reaching the estuarine area on the east of the province (Table 1). This is a curious record, given that this species was always found in the Subantarctic and west limit of the Patagonian subregions, very close to the Andes (Illies 1960a, McLellan 2001).

In Patagonia, as well as in many other regions of Argentina and the world over, mayflies and stoneflies are used as indicators of water quality and in biotic indexes (Miserendino & Pizzolón 1999, Pizzolón & Miserendino 2000). An adequate knowledge of the species distribution and information on their biotopes improves the

performance of biotic indexes. However, information about systematics, biology and ecology of these taxa in most of Patagonia is scarce. In this sense, collaboration between ecologists and systematists improves the knowledge of these taxa for use in water quality assessment and conservation of these species in Argentina.

Even though the present contribution is the result of a limited sampling effort, it increases by 100% the number of Ephemeroptera genera and by 65% the number of Plecoptera species recorded for Chubut province. In addition, it provides nine new country records for Argentina. These results further demonstrate the poor knowledge of the aquatic insect fauna in Patagonia and Chubut province. They also demonstrate the necessity of new and systematic entomological surveys in the area. Patagonian aquatic environments are endangered by the construction of hydroelectric dams, seeding of exotic fishes and by poor forestry practices. Long-term poor land use practices have resulted in severe desertification problems in broad areas in the steppe. All these activities threaten the biodiversity of these once pristine and fragile aquatic ecosystems (Pascual *et al.* 1998).

Acknowledgements

Our appreciation is extended to the Administración de Parques Nacionales and to Lic. Claudio Chehébar, Delegación Regional Patagonia, APN, for allowing the specimen collection within Los Alerces National Park, and Dr. Edward DeWalt and the anonymous reviewers for providing several suggestions that greatly improved the manuscript. This work was partially supported by the Argentinean Council of Scientific Research, grant number CONICET, PIP 5733. Special thanks to Dr. Miguel Archangelsky for helpful comments of the English version of the paper.

References

- Aubert, J. (1960) Contribution à l'étude des Notomemourinae (Plecoptera) de l'Amérique du Sud. *Bulletin de la Société Entomologique Suisse*, 33, 47–64.
- Claassen, P.W. (1936) New names for stoneflies (Plecoptera). *Annals of the Entomological Society of America*, 29, 622–623.
- Coronato, F.R. & Del Valle, H.F. (1988) Caracterización hídrica de las cuencas hidrográficas de la provincia del Chubut. Publicación Técnica. Cenpat-Conicet. Puerto Madryn. Chubut. Argentina, 183 pp.
- Demoulin, G. (1955a) Éphéméroptères nouveaux ou rares du Chili. *Bulletin de l'Institut Royal des Sciences Naturelles de Belgique*, 31, 1–15.
- Demoulin, G. (1955b) Éphéméroptères nouveaux ou rares du Chili. II. *Bulletin de l'Institut Royal des Sciences Naturelles de Belgique*, 31, 1–16.
- Demoulin, G. (1955c) Éphéméroptères nouveaux ou rares du Chili. III. *Bulletin de l'Institut Royal des Sciences Naturelles de Belgique*, 31, 1–30.
- Demoulin, G. (1966) Contribution à l'étude des éphéméroptères du Surinam. *Bulletin de l'Institut Royal des Sciences Naturelles de Belgique*, 42, 1–22.
- Domínguez, E. (1998) Ephemeroptera. In: Morrone, J. J. y Coscarón, S. (Eds.), *Biodiversidad de Artrópodos Argentinos. Una perspectiva biotaxonomica*. Ediciones Sur, La Plata, pp. 14–25.
- Domínguez, E., Hubbard M.D. & Pescador M.L. (1994) Los Ephemeroptera en Argentina. In: Castellanos, Z.A. (Ed.), *Fauna de Agua Dulce de la República Argentina*. FECIC, Buenos Aires, 33 (1), pp. 1–142.
- Domínguez, E., Hubbard, M.D., Pescador, M.L. & Molineri, C. (2001) Ephemeroptera. In: H.R. Fernández, E. Domínguez (Eds.), *Guía para la Determinación de los Artrópodos Bentónicos Sudamericanos*. EudeT, Tucumán, pp. 17–54.
- Eaton, A.E. (1882) An announcement of new genera of Ephemeridae. *Entomologist's Monthly Magazine*, 18, 207–208.
- Eaton, A.E. (1883–1888) A revisional monograph of Recent Ephemeridae or Mayflies. *Transactions of the Linnean Society of London, Zoology* (2) 3, 1–352, pl. 1–65.
- Edmunds, G.F. Jr. (1948) A new genus of mayflies from western North America (Leptophlebiinae). *Proceedings of the*

- Entomological Society of Washington*, 61, 141–146.
- Enderlein, G. (1909) Klassifikation der Plecopteren sowie Diagnosen neuer Gattungen und Arten. *Zoologischer Anzeiger*, 34, 385–419.
- Froehlich, C. G. (1960) Some gripopterygids and notonemurines (Plecoptera) from South America. *Lunds Universitets Arsskrift N.F.*, 56(13), 1–24.
- Gordon, N.D., Mc Mahon, T.A. & Finlayson, B.L. (1994) Stream Hydrology, an Introduction for Ecologists. Wiley & Sons, New York, 526 pp.
- Hauer, F.R. & Resh, V.H. 1996. Benthic macroinvertebrates. In: Hauer, F.R. & G.A. Lamberti (Eds), *Methods in Stream Ecology*. Academic Press, San Diego, California, pp. 339–369
- Hynes, H.B.N & Hynes, M.E. (1975) The life history of many of the stoneflies (Plecoptera) of south-eastern mainland Australia. *Australian Journal of Marine and Freshwater Research*, 27, 61–70.
- Hollmann, M.E.T. & Miserendino, M.L (2006) Habitat type and life history of the mayflies *Metamonius anceps* Eaton (Nesameletidae) and *Meridialaris chiloensis* Demoulin (Leptophlebiidae) in a Patagonian mountain stream. *Annales des Limnologie*, 42(4), 233–240.
- Hollmann, M.E.T. & Miserendino M.L. (2008) Life history and emergence patterns of stonefly species in mountain streams of the Futaleufú basin, Patagonia (Argentina). *Annales des Limnologie*, 44(2): 1-10.
- Illies, J. (1958) Abranchioperlidae, eine neue Plecopterenfamilie. *Investigaciones Zoológicas Chilenas*, 164, 26–41.
- Illies, J. (1960a) Penturoperlidae, eine neue Plecopterenfamilie. *Zoologischer Anzeiger*, 164, 26–41.
- Illies, J. (1960b) Archiperlaria, eine neue Unterordnung der Plecopteren. *Beiträge zur Entomologie*, 10, 661–697.
- Illies, J. (1961) Südamerikanische Notonemourinae und die Stellung der Unterfamilie im System der Plecopteren. *Mitteilungen der Schweizerischen Entomologischen Gesellschaft*, 34, 97–126.
- Illies, J. (1963) Revision der südamerikanische Gripopterygidae (Plecoptera). *Mitteilungen der Schweizerischen Entomologischen Gesellschaft*, 36, 145–248.
- Illies, J. (1964) Südamerikanische Perlidae (Plecoptera), besonders au Chile und Argentinien. *Beiträge zur Neotropischen Fauna*, 3, 207–233.
- Kluge, N. (1992) Cuban mayflies of the family Baetidae (Ephemeroptera). 2. Subgenera Caribaetis subgen. n. and Americabaetis subgen. n. of the genus Baetis s. l. *Zoologicheskij Zhurnal*, 4, 13–20.
- Klapálek, F. (1904) Plecopteren. In: *Hamburger Magalhaensische Sammelreise*, pp 1–30.
- Léon, R., Bran, D., Collantes, M., Paruelo, J.M. & Soriano, A. (1998) Grandes unidades de vegetación de la Patagonia extra andina. *Ecología Austral*, 8, 125–144.
- Lestage, J.A. (1931) Contribution à l'étude des Éphéméroptères. VIII. Les Éphéméroptères du Chili. *Bulletin et Annales de la Société Entomologique de Belgique*, 71, 41–60.
- Lugo-Ortiz, C.R. & McCafferty W.P. (1996) Taxonomy of the Neotropical genus *Americabaetis*, new status (Insecta: Ephemeroptera: Baetidae). *Studies on Neotropical Fauna and Environment*, 31, 156–169.
- Lugo-Ortiz, C.R. & McCafferty W.P. (1999) Three new genera of small minnow mayflies (Insecta: Ephemeroptera: Baetidae) from the Andes and Patagonia. *Studies on Neotropical Fauna and Environment*, 34, 88–104.
- Mabille, J. (1891) Sectio Plecoptera. In: *Néuroptères. Mission scientifique du Cap Horn, 1882–1883*, pp. 6–9.
- Mercado, M. & Elliot, S. (2004) Taxonomic revision of the genus *Metamonius* Eaton (Nesameletidae: Ephemeroptera), with notes on its biology and distribution. *Studies on Neotropical Fauna and Environment*, 39(2), 149–157.
- McLellan, I.D. (2001) A revision of the South American Austroperlidae (Plecoptera). *Aquatic Insects*, 23(3), 233–251.
- McLellan, I.D., Miserendino, M.L. & Hollmann, M.E.T. (2006) Two new species of *Notoperla* (Plecoptera: Gripopterygidae) and a redescription of *Notoperlopsis femina* Illies. *Zootaxa*, 1140, 63–68.
- McLellan, I.D., Wais I. & L.I. De Cabo (1990) The first record of stoneflies from the Malvinas/Falkland islands. *Aquatic Insects*, 12(3), 177–180.
- McLellan, I.D. & Zwick, P. (2007) New species of and keys to South American Gripopterygidae (Plecoptera). *Illiesia*, 3(4), 20–42.
- Miserendino, M.L. (1996) Las ninfas de plecópfera de ambientes lóticos del noroeste de Chubut. *Naturalia Patagónica. Serie Ciencias Biológicas*, 4(1–2), 75–83.
- Miserendino, M.L. (2001) Macroinvertebrate assemblages in Andean Patagonian rivers and streams. *Hydrobiologia*, 444, 147–158.
- Miserendino, M.L. (2006) Seasonal and spatial distribution of stoneflies in the Chubut river (Patagonia, Argentina). *Hydrobiologia*, 568, 263–274.
- Miserendino, M.L. & Hollmann, M.E.T. (2005) Patrones de emergencia de especies de Plecoptera en ríos del noroeste de Chubut (Patagonia-Argentina). *VI Congreso Argentino de Entomología*. 12 Septiembre. San Miguel de Tucumán. pp. 191.
- Miserendino, M.L. & Pizzolón, L.A. (1999) Rapid assessment of river water quality using macroinvertebrates: a family level biotic index for the Patagonic Andean zone. *Acta Limnologica Brasiliensia*, 11(2), 137–148.
- Miserendino, M.L. & Pizzolón, L.A. (2000) Macroinvertebrates of a fluvial system in Patagonia: altitudinal zonation and functional structure. *Archiv fur Hydrobiologie*, 150(1), 55–83.

- Miserendino, M.L. & Pizzolón, L.A. (2004) Interactive effects of basin features and land use change on macroinvertebrate communities of headwater streams in the Patagonian Andes. *River Research & Applications*, 20 (8), 967–982.
- Morrone, J.J. (2006) Biogeographic areas and transition zones of Latin America and the Caribbean Islands based on pan-biogeographic and cladistic analyses of the entomofauna. *Annual Reviews of Entomology*, 51, 467–494.
- Muzón, J. & Bachman, A.O. (1998) Plecoptera. In: Morrone, J. J. y Coscarón, S. (Eds.), *Biodiversidad de Artrópodos Argentinos. Una perspectiva biotaxonomica*. Ediciones Sur, La Plata, pp. 26–31.
- Muzón, J. & Pessacq, P. (2001) Nuevos registros de Ephemeroptera para la Meseta de Somuncurá, Río Negro. *Revista de la Sociedad Entomológica Argentina*, 60 (1–4), 67–78.
- Navás, L. (1918a) Insectos chilenos. *Boletín de la Sociedad Aragonesa de Ciencias Naturales*, 17, 212–230.
- Navás, L. (1918b). Algunos Insectos Neurópteros de la Argentina. *Phycis*, 4, 81–88.
- Navás, L. (1919) Insecta Nova. In: *Memoires della Pontificia Academia Romana Nuovi Lincei*, 2(5), 2–7
- Navás, L. (1928) Insectos neotrópicos. *Revista Chilena de Historia Natural*, 31, 231–232.
- Navás, L. (1929) Algunos Insectos de Chile. *Revista Chilena de Historia Natural*, 32, 327–329.
- Navás, L. (1930) Algunos Insectos de Chile. 3rd. Series. *Revista Chilena de Historia Natural*, 33, 326–334.
- Navás, L. (1933) Algunos Insectos de Chile. *Revista Chilena de Historia Natural*, 37, 231–232.
- Navás, L. (1934) Insectos suramericanos. *Revista de la Academia de Ciencias de Madrid*, 31, 155–184.
- Needham, J.G. & Murphy, H.E. (1924) Neotropical Mayflies. *Bulletin of the Loyd Library of Botany, Pharmacy and Materia Medica. 24. Entomological Series*, 4, 1–79.
- Nelson, C.H. (1973) Synopsis of the genus *Chilenoperla*. *The Pan Pacific Entomologist*, 49, 315–324.
- Nieto, C. (2004) South American Baetidae (Ephemeroptera): a New Generic Synonymy. *Studies on Neotropical Fauna and Environment*, 39(2), 95–101.
- Pascual, M.A., Orensanz, J.M., Parma, A.M. & Saba, S. (1998) The Patagonian Challenge: Melding Conservation with Development. Conservation Biology: Theory and Practice. In: Fiedler and Kareiva (Eds.), *Conservation Biology for the Coming Decade*, 2nd edition. Chapman and Hall, London, pp. 410–425.
- Pescador, M.L. & Peters, W.L. (1980) Two new genera of cool-adapted Leptophlebiidae (Ephemeroptera) from southern South America. *Annals of the Entomological Society of Americas*, 73, 332–338.
- Pescador, M.L. & Peters, W.L. (1982) Four new genera of Leptophlebiidae (Ephemeroptera: Atalophlebiinae). *Journal of the Kansas Entomological Society*, 58, 91–123.
- Pescador, M.L. & Peters, W.L. (1985) Biosystematics of the genus *Nousia* from Southern South America (Ephemeroptera: Leptophlebiidae: Atalophlebiinae). *Aquatic Insects*, 4, 1–19.
- Pescador, M.L. & Peters, W.L. (1991) Biosystematics of the genus *Penaphlebia* (Ephemeroptera: Leptophlebiidae: Atalophlebiinae) from South America. *Transactions of the American Entomological Society*, 117, 1–38.
- Pictet, F.J. (1841) *Historie Naturelle Générale et Particuliere des Insects Névroptères. Famille des Perlides*. Genève, 2 Bände.
- Pizzolón, L. & Miserendino, M.L. (2000) The performance of two regional biotic indices for running water quality in Northern Patagonian Andes. *Acta Limnologica Brasiliensia*, 13(1), 11–27.
- Richardson, J.S. (2001) Life cycle phenology of common detritivores from a temperate rainforest stream. *Hydrobiologia*, 455, 87–95.
- Scrimgeour, G.J. (1991) Life history and production of *Delatidium* (Ephemeroptera: Leptophlebiidae) in an unstable New Zealand river. *New Zealand Journal of Marine and Freshwater Research*, 25, 93–99.
- Traver, J.R. & Edmunds, G.F. Jr. (1968) A revision of the Baetidae with spatulate-clawed nymphs (Ephemeroptera). *Pacific Insects*, 10, 629–677.
- Ulmer, G. (1920) Neue Ephemeropteren. *Archiv für Naturgeschichte*, 85A (11), 1–80.
- Velásquez, S.M. & Miserendino, M.L. (2003) Habitat type and macroinvertebrate assemblages in low order Patagonian streams. *Archiv für Hydrobiologie*, 158(4), 461–483.
- Vera, A. (2006) Una nueva especie de Gripopterygidae de Chile, *Potamoperla testacea* n. sp. (Insecta: Plecoptera). *Gayana*, 70(2), 168–175.
- Wais, I.R., Villa, A.R. & Cabo, L.I. (1987) Las ninfas de plecópteros de los parques nacionales Nahuel Huapi y Lanin, Argentina. *Revista del Museo Argentino de Ciencias Naturales, Hidrobiología*, 6(3), 15–21.
- Zwick, P. (1972) On *Neonemura illiesi* nov. spec. and some other stoneflies from Chile (Ins. Plecoptera). *Studies on Neotropical Fauna*, 7, 95–100.