



Ephemeroptera (Insecta) Fauna of Northeast China

Yong-Tai Quan, Yeon Jae Bae¹⁾, Jong-Chul Jung²⁾,
and Jong-Wook Lee^{2)*}

Department of Biology, Harbin Normal University, Harbin, China

¹⁾Department of Biology, Seoul Women's University, Seoul 139-744, Korea

²⁾Department of Biology, Yeungnam University, Gyeongsan 712-749, Korea.

E-mail: jwlee1@ymail.ac.kr

Abstract A comprehensive literature review and examination of mayfly materials from Northeast China resulted in the recognition of 65 species, 28 genera, and 12 families of Ephemeroptera. Checklist, original reference sources, larval diagnoses, measurements, materials examined, distributions, and taxonomic remarks are provided. Macro-photos of larval body (dorsal view) and line-drawings of larval key characters are also provided.

Key words Mayflies, taxonomy, checklist

INTRODUCTION

Northeast China, commonly known as Manchuria, is located in 33°–41°N in latitude and 121°–135°E in longitude, and comprises Heilongjiang, Jilin, and Liaoning provinces. Ephemeroptera fauna from this area has been known by only several literatures. Hsu (1931–32, 1935–36) described two species of mayflies, and Imanishi (1940) reported 19 species of mayflies from this area. Gui (1985) included three species of mayflies from Jilin province in his "Catalogue of the Ephemeroptera of China." You and Gui (1995) included 10 species of mayfly adults from Northeast China. Bae (1997) presented a historical perspective of mayfly systematics in Northeast Asia. Bae and Liu (1999) reprinted 33 species of mayflies from Changbaishan area in Jilin province. Bae *et al.* (2000) reviewed the species of Northeast Asian Ephemeroptera in Imanishi's 1940 report. Lee *et al.* (2001) reported 30 species of mayflies (excluding 10 undetermined) from Heilongjiang and Jilin provinces.

The purpose of this study is to review and report mayfly fauna from Northeast China based on comprehensive literature review and materials examination.

MATERIALS AND METHODS

The materials used in this study include specimens of Seoul Women's University Aquatic Insect Collection, Yeungnam University Collection, and materials collected by the authors in 1999 and 2000 from Heilongjiang and Jilin provinces in Northeast China. Since this study has been concerned mainly on the faunistic survey of Ephemeroptera in the area, larval materials have been exclusively sampled and used, but other distributional records previously reported from this area are also included. Larvae were collected by hand nets and preserved in 80%

* To whom correspondence should be addressed.

Table 1. The abbreviations for larval body (mature) measurement used in this study

BL	: Body length
HL	: Head length
HW	: Head width
AL	: Antenna length
EL	: Longitudinal diameter of compound eye
EW	: Cross diameter of compound eye
DE	: Distance between compound eyes
PL	: Pronotum length
PW	: Pronotum width
CL	: Cerci length
TF	: Terminal filament length

Ethanol. The abbreviations used in larval body measurement are as in Table 1.

TAXONOMIC ACCOUNT

Family Ameletidae

1. *Ameletus costalis* (Matsumura) (Figs. 1, 83, 109, 140, 189)
Chimura costalis Matsumura, 1931: 1474.

Diagnosis. Body brown with pale markings. Labrum with V-shaped marking. Abdominal terga with triangular markings.

Measurement (mm). BL: 10.2, HW: 1.4, EL: 0.7, EW: 0.5, DE: 0.3, CL: 3.5.

Materials examined. Heilongjiang Heihe Sijiazi (5L, 26 VII 1997, YTQ), Yabuli Shishun (26L, 28 VII 1997, YTQ), Mudanjiangshi Liuchun (25L, 29 VII 1997, YTQ), Ninganxian Dongjingchengzhen (14L, 15 VII 2000). Jilin Yanjishi Xidong (2L, 3 VIII 2000, YTQ).

Distribution. China (Heilongjiang, Jilin), Japan, Korea, Russia.

2. *Ameletus montanus* Imanishi (Figs. 2, 110)
Ameletus montanus Imanishi, 1930: 265.

Diagnosis. Body brown. Outer canine of right mandible divided into three processes. Abdominal terga 3–7 with round markings. Cerci with broad band.

Measurement (mm). BL: 11.0, HL: 2.1, HW: 1.5, EL: 1.0, EW: 0.6, DE: 0.3, CL: 4.1.

Materials examined. Heilongjiang Mudanjiangshi Liuchun (2L, 29 VII 1997, YTQ). Jilin Antu Toudaobaihe (20L, 3 VI 1997, YJB), Antu Eradaobaihe (5L, 3 VI 1997, YJB), Antu Baihe (22L, 2 VI 1997, YJB), Antu Sandao (108L, 2 VI 1997, YJB), Helong (2L, 6 VI 1997, YJB), Helong Qingshanli (29L, 1 VI 1997, YJB), Helong Songxiaping (9L, 1 VI 1997, YJB).

Distribution. China (Heilongjiang, Jilin), Japan, Korea.

Family Baetidae

3. *Acentrella sibirica* (Kazlauskas) (Figs. 3, 82, 97, 126, 127, 159)
Pseudocloeon sibiricum Kazlauskas, 1963: 589.

Diagnosis. General body color brown. Vertex with irregular symmetrical brown markings. Maxillary palp 2-segmented; terminal segments apically round. Thorax relatively broad. Legs light brown; articulations brown; femora with single row of long setae along posterior margin; tibiae with single row of long setae dorsally; hindtibiae with additional row of short setae along posterior margin. Terga 1-8 with paired brown spots. Gills 1-7 oval. Terminal filament rudimentary.

Measurement (mm). ♂: BL: 3.8, HL: 0.7, HW: 0.92, AL: 0.22, EW: 0.17, DE: 0.62, CL: 2.8. ♀: BL: 3.7, HL: 0.6, HW: 0.8, AL: 1.0, EL: 3.8, EW: 0.17, DE: 0.45, CL: 2.7.

Materials examined. Heilongjiang Harbin Chenggaozi (4L, 4 VII 2000, YTQ), Ninganxian Dongjingchengzhen (3L, 15 VII 2000, YTQ).

Distribution. China (Heilongjiang), Japan, Korea, Russia.

4. *Baetis fuscatus* (Linnaeus) (Figs. 4, 128, 129, 160, 177)

Ephemera fuscata Linnaeus, 1761 No. 1477.

Diagnosis. Body brown. Maxillary palp segments 3 not clearly articulated. Terminal segments of labial palp with many hairlike setae. Abdominal terga 2-8 with paired opaque dark spots. Terminal filament well developed.

Measurement (mm). ♀: BL: 6.0, HL: 1.1, HW: 0.85, AL: 2.0, EL: 0.37, EW: 0.2, DE: 2.0, CL: 2.5.

Materials examined. Jilin Yanjishi Xiaowudao (2L, 27 VII 1997, YTQ), Yanji (63L, 6 VI 1997, YJB).

Distribution. China (Heilongjiang, Jilin, Liaoning), Japan, Korea, Mongolia, Russia.

5. *Baetis pseudothermicus* Kluge (Figs. 5, 56, 57, 84, 111, 112, 178)

Baetis pseudothermicus Kluge, 1983a: 53.

Diagnosis. Body light brown. Apical teeth of mandibles without denticle on outer margin. Maxillary palp with spinule on apex. Abdominal terga 3-4 and 6-8 darker. Gills 1-7 with distinct tracheae. Terminal filament well developed.

Measurement (mm). BL: 7.0, HL: 1.3, HW: 1.3, AL: 2.5, EL: 0.5, EW: 0.7, DE: 0.3, CL: 4.5, TF: 3.0.

Materials examined. Heilongjiang Daweniuhé (1L, 26 VII 1997, YTQ), Hailinshi Pinghe (4L, 28 VII 1997, YTQ), Harbin (3L, 29 VII 1997, YTQ), Harbin Chenggaozi (7L, 4 VII 2000, YTQ), Ninganxian Dongjingchengzhen (23L, 15 VII 2000, YTQ). Jilin Yanjishi Xidong (64L, 3 VIII 2000, YTQ) Antu Toudaobaihe (46L, 3 VI 1997, YJB), Antu Eradaobaihe (88L, 3 VI 1997, YJB), Antu Baihe (16L, 2 VI 1997, YJB), Antu Sandao (163L, 1 VI 1997, YJB), Helong (1L, 6 VI 1997, YJB), Wolong Xianfeng (100L, 6 VI 1997, YJB), Helong Qingshanli (12L, 1 VI 1997, YJB).

Distribution. China (Heilongjiang, Jilin), Korea, Russia.

6. *Baetiella tuberculata* (Kazlauskas) (Fig. 6)

Pseudocloeon (Baetiella) tuberculatum Kazlauskas, 1963: 583.

Diagnosis. Body brown. Vertex with 3-4 pairs of symmetrical brown markings. Maxillary palp 2-segmented. Pronotum with 2 pairs of dark brown markings medially. Legs articulations brown. Claws brown with denticles and paired long terminal setae. Terminal filament rudimentary.

Measurement (mm). ♂: BL: 4.25, HL: 0.8, HW: 0.9, AL: 1.5, EL: 0.5, EW: 0.35, DE: 0.25,

CL: 4.1. ♀: BL: 5, HL: 0.7, HW: 0.8, EL: 0.3, EW: 0.2, DE: 0.38, CL: 4.5.

Materials examined. **Heilongjiang** Hailinshi Sanbuluo (27L, 28 VII 1997, YTQ), Shangzhishi Maorshanzhen (5L, 4 VI 1999, YTQ), Mudanjiangshi Liuchun (17L, 29 VII 1997, YTQ), Ninganxian Dongjingchengzhen (36L, 15 VII 2000, YTQ). **Jilin** Yanjishi Xiaowudao (6L, 27 VII 2000; 12L, 30 VII 2000, YTQ).

Distribution. China (Heilongjiang, Jilin), Korea, Russia.

7. *Cloeon dipterum* (Linnaeus) (Figs. 7, 58, 183)

Ephemera diptera Linnaeus, 1761, No. 1477.

Diagnosis. Body brown. Claws with two rows of well developed denticles. Gills 1–6 paired; gills 7 single. Posterior margin of abdominal segment 10 round.

Measurement (mm). BL: 4.1, HL: 1.1, HW: 3.4, AL: 2.0, EL: 0.35, EW: 0.32, DE: 0.3, CL: 4.1.

Materials examined. **Heilongjiang** Mudanjiangshi Liuchun (2L, 29 VII 1997, YTQ).

Distribution. China (Heilongjiang, Jilin, Liaoning), Japan, Korea, Mongolia, Russia.

8. *Procloeon pennulum* (Eaton) (Figs. 8, 60, 64, 85, 113, 162, 208)

Centroptilum pennulum Eaton, 1870: 2.

Diagnosis. Body brown. Claws with two rows of rudimentary denticles basally, with terminal hair. Gills 1–6 paired; gills 7 single. Posterior margin of abdominal segment 10 straight.

Measurement (mm). BL: 7.5, HL: 1.1, HW: 1.0, AL: 5.0, EL: 0.32, EW: 0.25, DE: 0.2, CL: 2.5.

Materials examined. **Heilongjiang** Hailinshi Pinghe (3L, 28 VII 1997, YTQ). **Jilin** Yanji (3L, 6 VI 1997, YJB).

Distribution. China (Heilongjiang, Jilin), Korea, Russia.

Family Caenidae

9. *Brachycercus* YUa (Figs. 9, 114, 141, 184, 190)

Diagnosis. Body light yellow. Head with three projections. Prothorax with sublateral tubercles. Legs slender, with hair-like setae. Abdominal segment 2–9 with posterolateral projections; posterolateral projections 8–9 distinct. Caudal filaments light yellow, with whorls of setae at end of each segment.

Measurement (mm). BL: 6.4, HW: 1.2, AL: 1.5, EL: 0.35, EW: 0.32, DE: 0.8, CL: 3.2, TF: 3.8.

Materials examined. **Heilongjiang** Sijiazi (1L, 26 VII 1997, YTQ), Shangzhishi Maorshanzhen (1L, 4 VI 1999, YTQ), Harbin (1L, 29 VII 1999, YTQ).

Distribution. China (Heilongjiang).

Remarks. This species is distinguished from *Brachycercus tubulatus* Tshernova by the pronotal tubercles and posterolateral projections on the abdominal segments 8–9.

10. *Caenis nishinoae* Malzacher

Caenis nb: Imanishi, 1940: 212.

Caenis nishinoae Malzacher, 1996: 1.

Diagnosis. Body yellowish brown, with dark brown markings. Head with W-shaped black

marking. Legs light, with yellowish brown bands; claws simple.

Measurement (mm). BL: 4. HL: 0.7, HW: 0.85, AL: 2, EW: 0.25, DE: 0.45, CL: 3.7.

Materials examined. **Jilin** Yanjishi Yanxiaoqiao (1L, 2 VIII 2000, YTQ), Yanjishi Xidong (2L, 3 VIII 2000, YTQ).

Distribution. China (Jilin, Liaoning), Korea, Japan.

Remarks. Imanishi (1940) reported *Caenis* nb from Liaoning province. *Caenis* nb was later associated with *Caenis nishinoae* Malzacher by Hwang and Bae (1999).

11. *Caenis na* (Figs. 10, 138, 139, 161, 191)

Caenis na: Imanishi, 1940: 211.

Diagnosis. Body brown, with tiny light spots throughout dorsal body. Head with web-like markings. Basal and second antennal segments dark brown. Pronotum with large and acute anterolateral processes. Legs and cerci with brown band.

Measurement (mm). BL: 6.4, HL: 1.0, HW: 1.4, AL: 3.0, EL: 0.35, EW: 0.27, DE: 1.0, CL: 6.5.

Materials examined. **Heilongjiang** Sijiazi (5L, 26 VII 1997, YTQ), Harbin (7L, 29 VII 1997, YTQ). **Jilin** Yanjishi Yanxiaoqiao (5L, 2 VIII 2000, YTQ), Yanjishi Xidong (41L, 3 VIII 2000, YTQ).

Distribution. China (Liaoning), Korea.

12. *Caenis nc*

Caenis nc: Imanishi, 1940: 214.

Materials examined. Not available.

Locality. **Heilongjiang** Hailaerhaishuihe (Imanishi, 1940).

Distribution. China (Heilongjiang).

Family Ephemerellidae

13. *Cincticostella levanidovae* (Tshernova) (Figs. 11, 52, 63, 62, 86, 115, 116, 142, 163) *Ephemerella levanidovae* Tshernova, 1952: 274.

Diagnosis. Body brown. Head without occipital tubercles; genae truncate. Maxillae with well developed palp. Thoracic nota without tubercles or ridges. Abdominal terga 5-8 with well developed tubercles. Caudal filaments with stout setae on every segment.

Measurement (mm). BL: 9.6, HL: 1.5, HW: 1.7, AL: 2.25, EL: 1, EW: 0.8, DE: 1.1, CL: 6.5.

Materials examined. **Heilongjiang** Heile Xingfuxiang (5L, 26 VII 1997, YTQ), Sijiazi (10L, 26 VII 1997, YTQ), Hailinshi Pinghe (4L, 28 VII 1997, YTQ), Hailinshi Sanbuluo (7L, 28 VII 1997, YTQ), Hailinshi Shunqiao (26L, 28 VII 1997, YTQ), Yanshouxian Yanhe (5L, 6 VI 1999, YTQ), Mudanjiangshi Liuchun (11L, 29 VII 1997, YTQ), Ninganxian Dongjingchengzhen (15L, 15 VII 2000, YTQ). **Jilin** Yanjishi Xiaowudao (14L, 30 VI 2000, YTQ), Yanjishi Xidong (41L, 3 VIII 2000, YTQ), Antu Toudaobaihe (12L, 3 VI 1997, YJB), Antu Sandao (10L, 1 VI 1997, YJB), Antu Sidaobaihe (1L, 1 VI 1997, YJB), Wolong (1L, 6 VI 1997, YJB).

Distribution. China (Heilongjiang, Jilin), Japan, Korea, Russia.

14. *Cincticostella tshernovae* Bajkova (Figs. 12, 59, 65, 66, 117, 143, 164, 185, 192, 193) *Cincticostella tshernovae* Bajkova, 1962: 203.

Diagnosis. Body reddish brown to dark brown with purple to blackish purple markings. Abdominal terga 5–9 with large paired submedian tubercles, with short heavy setae. Caudal filaments less than one-half body length.

Measurement (mm). BL: 10.3, HL: 2.0, HW: 2.4, AL: 2.7, EL: 0.65, EW: 0.5, DE: 1.5, CL: 5.0.

Materials examined. Jilin Yanjishi Xiaowudao (3L, 30 VI 2000, YTQ), Hailinshi Pinghe (2L, 28 VII 1997, YTQ).

Distribution. China (Jilin), Japan, Korea, Russia.

15. *Drunella aculea* (Allen)

Ephemerella basalis Na: Imanishi, 1940: 196.

Materials examined. Not available.

Locality. Jilin Antu Eradaobaihe (Bae and Liu, 1999), Antu baihe (Bae and Liu, 1999), Antu Sandao (Bae and Liu, 1999), Helong, Wolong (Bae and Liu, 1999), Helong Songxiaping (Bae and Liu, 1999).

Distribution. China (Jilin), Korea, Russia.

16. *Drunella cryptomeria* (Imanishi) (Figs. 13, 119, 194)

Ephemerella cryptomeria Imanishi, 1937a: 328.

Diagnosis. Body brown. Head with three long clypeal tubercles. Frontal clypeal edge notched. Forefemora dosally with distinct ridges. Tarsal claw with three denticles. Abdominal terga 2–7 with paired pale projection. Caudal filaments with whorls of setae at end of each segment.

Measurement (mm). ♂: BL: 8.5, HL: 1.3, HW: 0.9, AL: 1.2, EL: 0.7, EW: 0.6, DE: 1.7, CL: 2.5, TF: 2.7. ♀: BL: 10.8, HL 1.5, HW: 2.1, EL: 0.7, EW: 0.5, DE: 1.0, CL: 5.5, TF: 5.8.

Materials examined. Heilongjiang Hailinshi Sanbuluo (11L, 28 VII 1997, YTQ), Hailinshi Shunqiao (4L, 28 VII 1997, YTQ). Jilin Yanjishi Xiaowudao (7L, 30 VII 2000, YTQ), Antu Baihe (7L, 2 VI 1997, YJB), Antu Sandao (12L, 1 VI 1997, YJB), Helong (2L, 6 VI 1997, YJB), Wolong (1L, 6 VI 1997, YJB), Helong Songxiaping (32L, 1 VI 1997, YJB), Helong Songxiaping (8L, 1 VI 1997, YJB).

Distribution. China (Heilongjiang, Jilin), Japan, Korea, Russia.

17. *Drunella fusongensis* Su and Gui

Drunella fusongensis Su and Gui, 1995: 451.

Materials examined. Not available.

Locality. Jilin Fusongxian (Su and Gui, 1995).

Distribution. China (Jilin).

Remarks. Su and Gui (1995) described adult of this species from Jilin province.

18. *Drunella lepnevae* (Tshernova) (Figs. 14, 67, 121, 122, 144)

Ephemerella lepnevae Tshernova, 1949: 152.

Diagnosis. Body brown or light brown. Head without clypeal tubercles, with paired occipital tubercles. Anterior margin of forefemora with small denticles. Abdominal terga 2–9 with paired pale tubercle. Caudal filaments with row of dense dorsal setae and whorls of setae at each segment.

Measurement (mm). BL: 9.0, HL: 2.1, HW: 2.2, AL: 1.5, EL: 1.1, EW: 0.8, DE: 0.85, CL: 8.0, TF: 9.0.

Materials examined. **Heilongjiang** Hailinshi Sanbuluo (3L, 28 VII 1997, YTQ), Jilin Antu Toudaobaihe (3L, 3 VI 1997, YJB), Antu Eradaobaihe (18L, 3 VI 1997, YJB), Antu Baihe (261L, 2 VI 1997, YJB), Antu Sandao (2L, 1 VI 1997, YJB), Helong Songxiaping (1L, 1 VI 1997, YJB).

Distribution. China (Heilongjiang, Jilin), Korea, Mongolia, Russia.

19. *Drunella solida* (Bajkova) (Figs. 15, 45, 68, 100, 118, 145, 165, 195, 196)
Ephemerella solida Bajkova, 1980: 74.

Diagnosis. Body brown, with distinct light markings. Head with three frontal tubercles; clypeal margin round. Forefemora with frontal spines, without dorsal ridges; tibiae and tarsi with one broad transverse stripe.

Measurement (mm). BL: 12.4, HL: 1.6, HW: 2.6, EL: 0.8, EW: 0.7, DE: 1.1, CL: 8.0.

Materials examined. Jilin Yanjishi Xiaowudao (1L, 27 VI 1999, YTQ), Antu Sandao (1L, 1 VI 1997, YJB).

Distribution. China (Jilin), Korea, Russia.

20. *Drunella triacantha* (Tshernova)

Ephemerella trispina na: Imanishi, 1940: 193.
Ephemerella triacantha Tshernova, 1949: 151.

Materials examined. Not available.

Locality. Jilin Antu Baihe (Bae and Liu, 1999).

Distribution. China (Jilin), Japan, Korea, Mongolia, Russia.

21. *Ephacerella longicaudata* (Uéno)

Ephemerella longicaudata Uéno, 1928: 42.
Ephacerella longicaudata (Uéno): Pacif., 1994: 283.

Materials examined. Not available.

Locality. **Heilongjiang** (You and Gui, 1995).

Distribution. China (Heilongjiang, Jilin, Liaoning), Japan, Korea, Russia.

22. *Ephemerella antuensis* Su and You

Ephemerella (Ephemerella) antuensis Su and You, 1989: 181.

Materials examined. Not available.

Locality. Jilin Antuxian (You and Gui, 1995).

Distribution. China (Heilongjiang, Jilin).

Remarks. Su and You (1989) described adult of this species from Jilin province.

23. *Ephemerella aurivillii* (Bengtsson) (Fig. 16)

Chitonophora aurivillii Bengtsson, 1909: 8.

Diagnosis. Body color brown or dark brown. Head without frontal tubercles. Abdominal terga without tubercle, with light longitudinal stripes. Caudal filaments with whorls of setae and lateral setae.

Measurement (mm). BL: 12.0, HL: 1.8, HW: 2.0, AL: 3, EL: 0.5, EW: 0.4, DE: 1.2, CL: 7.5.

Materials examined. Jilin Antu Eradaobaihe (5L, 3 VI 1997, YJB), Antu Baihe (10L, 2 VI 1997, YJB), Antu Sandao (5L, 1 VI 1997, YJB), Helong (18L, 6 VI 1997, YJB), Wolong Xianfeng (1L, 6 VI 1997, YJB), Helong Qingshanli (20L, 1 VI 1997, YJB), Helong Songxiaping (2L, 1 VI 1997, YJB).

Distribution. China (Jilin), Japan, Korea, Mongolia, Russia.

24. *Ephemerella changbaishanensis* Su and You

Ephemerella changbaishanensis Su and You, 1988: 63.

Materials examined. Not available.

Locality. Jilin Fushongxian (You and Gui, 1995).

Distribution. China (Jilin).

Remarks. Su and You (1988) described adult of this species from Jilin province.

25. *Ephemerella dentata* Bajkova (Figs. 17, 53, 69, 87, 101, 197)

Ephemerella dentata Bajkova, 1967: 331.

Diagnosis. Body light brown. Head without frontal tubercles. Forefemora with subapical row of stout setae. Abdominal terga with moderately developed submedian tubercles. Caudal filaments with whorls of setae and lateral setae.

Measurement (mm). ♂: BL: 6.0, HL: 0.8, HW: 1.4, AL: 3.5, EL: 0.5, EW: 0.7, DE: 0.2, CL: 4.5, TF: 4.8. ♀: BL: 6.7, HL: 1.2, HW: 1.6, AL: 2.1, EL: 0.5, EW: 0.3, DE: 0.8, CL: 4.7.

Materials examined. Heilongjiang Hulan (13L, 23 VII 1999, YTQ), Yabuli Shishun (8L, 28 VII 1997, YTQ), Hailinshi Pinghe (61L, 28 VII 1997, YTQ), Shangzhishi Maorshanzhen (60L, 4 VI 1999, YTQ), Mudanjiangshi Liuchun (25L, 29 VII 1997, YTQ). Jilin Yanjishi Xiaowudao (32L, 27 VI 1999, YTQ), Yanjishi Yanxiqiao (4L, 2 VIII 2000, YTQ), Antu Toudaobaihe (2L, 3 VI 1997, YJB), Antu Baihe (11L, 2 VI 1997, YJB), Antu Sandao (234L, 1 VI 1997, YJB), Antu Sidaobaihe (7L, 1 VI 1997, YJB), Helong (1L, 6 VI 1997, YJB), Wolong (25L, 6 VI 1997, YJB), Helong Songxiaping (138L, 1 VI 1997, YJB), Yanji (1L, 6 VI 1997, YJB).

Distribution. China (Heilongjiang, Jilin), Russia.

26. *Ephemerella fusongensis* Su and You

Ephemerella fusongensis Su and You, 1988: 64.

Materials examined. Not available.

Locality. Jilin Fushongxian (You and Gui, 1995).

Distribution. China (Jilin).

Remarks. Su and You (1988) described adult of this species from Jilin province.

27. *Ephemerella kozhovi* Bajkova (Figs. 18, 120, 146, 166, 198, 199, 200)

Ephemerella kozhovi Bajkova, 1967: 327.

Diagnosis. Body brown with broad traverse bands across basal part of wingpads and on abdominal terga 5–6. Head and thorax without distinct humps. Abdominal tergum 2–9 with rudimentary tubercles. Caudal filaments with whorls of setae and lateral setae.

Measurement (mm). ♂: BL: 7.5, HL: 1.3, HW: 1.6, AL: 1.7, EL: 0.75, EW: 0.7, DE: 0.65.

CL: 3.8. ♀: BL: 7.6, HL: 1.4, HW: 1.5, AL: 1.8, EL: 0.55, EW: 0.35, DE: 0.8, CL: 4.0.
Materials examined. Heilongjiang Yabuli Shishun (4L, 28 VII 1997, YTQ). Jilin Antu Eradaobaihe (2L, 3 VI 1997, YJB), Antu Baihe (8L, 2 VI 1997, YJB).
Distribution. China (Heilongjiang), Korea, Russia.

28. *Serratella setigera* (Bajkova) (Fig. 19)

Ephemerella setigera Bajkova, 1967: 333.

Diagnosis. Body brown with dark markings. Head and thorax without distinct tubercles. Abdominal terga 3–9 with paired tubercles. Caudal filaments with whorls of setae, without lateral setae, with dark bands on alternate segments.

Measurement (mm). BL: 6.0, HL: 0.9, HW: 1.25, AL: 1.5, EL: 0.5, EW: 0.4, DE: 0.5, CL: 3.1.

Materials examined. Heilongjiang Hailinshi Pinghe (9L, 28 VII 1997, YTQ), Mudanjiangshi Liuchun (6L, 29 VII 1997, YTQ). Jilin Yanjishi Xiaowudao (3L, 30 VI 2000).

Distribution. China (Heilongjiang, Jilin), Russia.

29. *Serratella zapekinae* (Bajkova) (Figs 20, 70, 88, 123, 186, 201)

Ephemerella zapekinae Bajkova, 1967: 329.

Diagnosis. Body brown. Head and thorax with dark tiny marking. Abdominal terga 3–8 with paired tubercles. Caudal filaments with whorls of setae, without lateral setae, without dark bands.

Measurement (mm). ♂: BL: 6.2, HL: 1.3, HW: 1.6, AL: 2.0, EL: 0.7, EW: 0.65, DE: 0.6. CL: 5.2. ♀: BL: 5.75, HL: 1.2, HW: 12.5, AL 1.9, EL: 0.4, EW: 0.25, DE: 0.8, CL: 5.0.

Materials examined. Heilongjiang Sijiazi (2L, 26 VII 1997, YTQ).

Distribution. China (Heilongjiang), Russia.

30. *Uracanthella rufa* (Imanishi) (Figs 21, 71, 102, 124, 125, 147, 167, 187)

Ephemerella rufa Imanishi, 1937a: 327.

Diagnosis. Body brown with light paired longitudinal stripes along dorsal body. Mandibles large. Abdominal terga without tubercles. Caudal filaments with whorls of setae and lateral setae.

Measurement (mm). ♂: BL: 5.9, HL: 1.45, HW: 1.4, CL: 3.2. ♀: BL: 6.0, CL: 3.0, HW: 1.35, HL: 1.4, AL: 1.5, EW: 0.2, EL: 0.4, DE: 0.9.

Materials examined. Heilongjiang Heihe Xingfuxiang (37L, 26 VII 1997, YTQ), Hailinshi Pinghe (3L, 28 VII 1997, YTQ), Shangzhi Maorshanzen (21L, 4 VI 1999, YTQ), Ning'anxian Dongjingchengzhen (31L, 15 VII 2000, YTQ).

Distribution. China (Heilongjiang), Korea, Mongolia, Russia.

Family Ephemeridae

31. *Ephemera kirinensis* Hsu

Ephemera kirinensis Hsu, 1931: 40.

Materials examined. Not available.

Locality. Jilin (You and Gui, 1995).

Distribution. China (Jilin).

Remarks. Hsu (1931) described adult of this species from Jilin province.

32. *Ephemera orientalis* McLachlan (Fig. 22)

Ephemera orientalis McLachlan, 1875: 167.

Diagnosis. Body color light yellow to light brown. Abdominal tergum 7–9 with paired three-lined stripes.

Measurement (mm). ♂: BL: 18.0, HL: 1.8, HW: 1.7, AL: 5.5, EL: 0.75, EW: 0.7, DE: 0.95, PL: 3.2, PW: 4.3, CL: 5.4 TF: 5.7.

Materials examined. Heilongjiang Hulan (1L, 23 VII 1999, YTQ), Ninganxian Dongjingchengzhen (5L, 15 VII 2000, YTQ). Jilin Yanjishi Yanxiaoqiao (1L, 2 VIII 2000, YTQ).

Distribution. China (Heilongjiang, Jilin, Liaoning), Japan, Korea, Mongolia, Russia.

33. *Ephemera sachalinensis* Matsumura

Ephemera sachalinensis Matsumura, 1911: 9.

Materials examined. Not available.

Locality. Heilongjiang Xiaoxinganling (You and Gui, 1995).

Distribution. China (Heilongjiang, Jilin, Liaoning), Korea, Russia.

Remarks. You and Gui (1995) reported adult of this species from Heilongjiang province.

34. *Ephemera separigata* Bae (Fig. 23)

Ephemera separigata Bae, 1995: 160.

Diagnosis. Body color light yellow to light brown. Abdominal tergum 7–9 with relatively narrow lateral stripes.

Measurement (mm). BL: 21.0, HL: 2.6, HW: 2.6, AL: 6.7, EL: 0.8, EW: 0.9, DE: 1.6, CL: 10

Materials examined. Jilin Antu Baihe (10L, VI 1997, YJB), Antu Sandao (10L, VI 1997, YJB).

Distribution. China (Jilin), Korea.

35. *Ephemera strigata* Eaton (Figs. 24, 72, 148, 168, 209)

Ephemera strigata Eaton, 1892: 302.

Diagnosis. Body light yellow or light brown. Abdominal tergum 7–9 with relatively broad sublateral stripes.

Measurement (mm). ♂: BL: 6.5, HL: 0.6, HW: 0.65, AL: 3.7, EL: 0.4, EW: 0.5, DE: 0.7, CL: 2.2. ♀: BL: 9.7, HL: 1.0, HW: 1.1, EL: 0.6, EW: 0.7, DE: 0.85, CL: 3.5, TF: 4.5.

Materials examined. Heilongjiang Sijiazi (1L, 26 VII 1997, YTQ), Hailinshi Pinghe (1L, 28 VII 1997, YTQ). Jilin Antu Baihe (3L, VI 1997 YJB).

Distribution. China (Heilongjiang, Jilin), Japan, Korea, Mongolia, Russia.

Family Heptageniidae

36. *Cinygmula brunnea* Tiunova (Fig. 25)

Cinygmula brunnea Tiunova, 1990: 814.

Diagnosis. Body brown. Head dark brown. Legs yellowish brown, with dark markings.

Abdominal terga 2–5 with light paired markings. Caudal filaments brown, basally somewhat lighter.

Measurement (mm). BL: 13.0, HL: 2.2, HW: 3.2, AL: 3, EL: 1.2, EW: 0.8, DE: 1.1, CL: 11.0.

Materials examined. Jilin Helong Qingshanli (25L, VI 1997, YJB), Wolong Xianfeng (10L, 6 VI 1997, YJB), Antu Eradaobaihe (4L, 3 VI 1997, YJB), Antu Sandao (95L, VI 1997, YJB) Antu Baihe (3L, VI 1997, YJB), Antu Sandao (10L, VI 1997, YJB), Wolong Xianfeng (16L, 6 VI 1997, YJB), Helong Qingshanli (25L, VI 1997, YJB).

Distribution. China (Jilin), Russia.

37. *Cinygmula hirasana* (Imanishi) (Figs. 26, 46, 54, 73)

Cinygma hirasana Imanishi, 1935: 214.

Diagnosis. Body brown. Thorax with distinct median light line and two light small spots on each side of the scutellum. Femora apically with light longitudinal markings. Abdominal terga 2–9 with light triangular markings.

Measurement (mm). BL: 9.2, HL: 2.0, HW: 2.6, AL: 2.5, EL: 0.8, EW: 0.6, DE: 0.8, CL: 9.0.

Materials examined. Heilongjiang Heihe Xingfuxiang (5L, 26 VII 1997, YTQ), Heihe Jinhengnongchang (3L, 26 VII 1997, YTQ), Sijiazi (2L, 26 VII 1997, YTQ), Shangzhishi Maorshanzen (8L, 4 VI 1999, YTQ). Jilin Yanjishi Xiaowudao (12L, 27 VII 2000, YTQ), Yanjishi Xiaowudao (20L, 27 VII 2000, YTQ), Yanjishi Xiaowudao (42L, 30 VI 2000, YTQ), Antu Toudaobaihe (3L, 3 VI 1997, YJB), Antu Eradaobaihe (35L, 3 VI 1997, YJB), Antu Baihe (29L, VI 1997, YJB), Antu Sandao (108L, VI 1997, YJB), Wolong Xianfeng (35L, 6 VI 1997, YJB), Wolong (11L, 6 VI 1997, YJB), Helong Qingshanli (25L, VI 1997, YJB), Helong Songxiaping (7L, VI 1997, YJB).

Distribution. China (Heilongjiang, Jilin), Japan, Korea, Russia.

38. *Cinygmula kurenzovi* (Bajkova) (Fig. 27)

Cinygma kurenzovi Bajkova, 1965: 301.

Diagnosis. Body color light yellowish brown, with brown markings. Labrum semicircular, reddish brown. Pronotum lateral margin hyaline, with Z-shape light stripes. Abdominal terga 2–9 with pair light spots. Gills white, with dark brown marking.

Measurement (mm). BL: 9.5, HL: 1.4, HW: 2.2, AL: 1.5, EL: 0.7, EW: 0.5, DE: 0.7, CL: 7.0, TF: 8.0.

Materials examined. Jilin Antu Toudaobaihe (2L, 3 VI 1997, YJB), Antu Eradaobaihe (18L, 3 VI 1997, YJB), Antu Baihe (15L, VI 1997, YJB), Antu Sandao (50L, VI 1997, YJB), Wolong Xianfeng (18L, 6 VI 1997, YJB), Wolong (6L, 6 VI 1997, YJB), Helong Qingshanli (13L, VI 1997, YJB), Helong Songxiaping (4L, VI 1997, YJB).

Distribution. China (Heilongjiang), Korea, Russia.

39. *Cinygmula levanidovi* Tshernova and Belov (Fig. 28)

Cinygma levanidovi Tshernova and Belov, 1982: 193.

Diagnosis. Body color light brown with light or brown markings. Head yellowish brown; middle of anterior margin with a pit. Pronotum light yellowish brown, with median longitudinal stripes. Forelegs with light yellowish brown markings; posterior margin with hairlike setae. Gills white.

Measurement (mm). BL: 7.5, HL: 1.8, HW: 2.0, AL: 1.5, EL: 0.7, EW: 0.4, DE: 0.7, CL: 7.0, TF: 7.5.

Materials examined. **Jilin** Antu Toudaobaihe (1L, 3 VI 1997, YJB), Antu Eradaobaihe (9L, 3 VI 1997, YJB), Antu Baihe (8L, VI 1997, YJB), Antu Sandao (28L, VI 1997, YJB), Wolong Xianfeng (9L, 6 VI 1997, YJB), Wolong (3L, 6 VI 1997, YJB), Helong Qingshanli (7L, VI 1997, YJB), Helong Songxiaping (2L, VI 1997, YJB).

Distribution. China (Jilin), Russia.

40. *Ecdyonurus dracon* Kluge (Figs. 29, 74, 89, 103, 130, 131, 150, 169, 179)
Ecdyonurus dracon Kluge. 1983b: 27.

Diagnosis. Body brown to dark brown. Frontal area of head without marking. Abdominal terga 4–5, 8–9 with U-shaped marking. Caudal filaments with whorls of spines at apex of each segment.

Measurement (mm). BL: 12.9, HL: 2.3, HW: 3.6, AL: 2.5, EL: 1.15, EW: 1.1, DE: 1.2.

Materials examined. **Heilongjiang** Ninganxian Dongjingchengzhen (2L, 15 VII 2000, YTQ).

Distribution. China (Heilongjiang), Korea, Russia.

41. *Ecdyonurus kibunensis* Imanishi (Figs. 30, 47, 75, 151, 170)
Ecdyonurus kibunensis Imanishi. 1936: 540.

Diagnosis. Body brown. Head with two obscure light spots on middle portion of frontal margin. Legs yellow; femora with brownish spot. Abdominal terga 4–5 with U-shaped marking.

Measurement (mm). BL: 5.4, HL: 1.4, HW: 2.0, EL: 0.7, EW: 0.6, DE: 0.6, CL: 7.5.

Materials examined. **Jilin** Yanjishi Xiaowudao (6L, 27 VII 2000, YTQ), Yanjishi Xiaowudao (2L, 30 VI 2000, YTQ), Yanjishi Yanxiaoqiao (12L, 2 VIII 2000, YTQ), Antu Toudaobaihe (1L, 3 VI 1997, YJB), Antu Baihe (3L, VI 1997, YJB).

Distribution. China (Jilin), Korea, Russia, Japan, Mongolia.

42. *Ecdyonurus levis* (Navás) (Figs. 31, 132, 152, 171, 180)
Epeorus levis Navás. 1912: 415.

Diagnosis. Body brown with light markings. Frontal margin of head with two paired round markings. Abdominal terga with distinct light markings. Caudal filaments with whorls of setae at apex of each segment.

Measurement (mm). ♂: BL: 9.2, HL: 1.9, HW: 2.95, EL: 1.1, EW: 1.4, DE: 0.1, CL: 12. ♀: BL: 11.6, HL: 2.3, HW: 3.4, EL: 1.2, EW: 1.2, DE: 0.8, TF: 15.0.

Materials examined. **Heilongjiang** Hailinshi Shunqiao (12L, 28 VII 1997, YTQ), Ninganxian Dongjingchengzhen (9L, 15 VII 2000, YTQ), **Jilin** Yanjishi Xiaowudao (4L, 27 VII 2000, YTQ), Yanjishi Yanxiaoqiao (13L, 2 VIII 2000, YTQ).

Distribution. China (Heilongjiang, Jilin), Korea, Russia.

43. *Ecdyonurus yoshidae* Takahashi
Ecdyonurus yoshidae Takahashi. 1924: 380.

Materials examined. Not available.

Locality. **Heilongjiang** Mudanjiangjingpohu (Imanishi, 1940). Liaoning Dalian (Imanishi,

1940), Shenyangtaizihe (Imanishi, 1940).

Distribution. China (Heilongjiang, Liaoning, Neimenggu), Korea, Russia, Japan.

44. *Epeorus curvatus* Matsumura (Figs. 32, 76, 92, 104, 153)

Epeorus curvatus Matsumura, 1931: 1477.

Diagnosis. Body brown. Frontal margin of head with paired C-shaped markings. Each tibia with brownish spot. Abdominal terga 2–10 with brownish median line. Abdominal segment 2–7 with two lateral projections. Tails two.

Measurement (mm). ♂: BL: 9.8, HL: 2.0, HW: 2.7, AL: 2.5, EL: 1.0, EW: 0.7, DE: 0.4, CL: 15.6. ♀: BL: 9.5, HL: 2.1, HW: 2.8, AL: 2.7, EL: 0.9, EW: 0.6, DE: 0.8, CL: 13.5.

Materials examined. Jilin Yanjishi Xiaowudao (1L, 27 VII 2000, YTQ), Antu Baihe (1L, VI 1997, YJB).

Distribution. China (Heilongjiang, Jilin, Liaoning), Korea, Russia, Japan.

45. *Epeorus latifolium* Uéno

Epeorus latifolium Uéno, 1928: 34.

Materials examined. Not available.

Locality. Liaoning Wangyemiao (Imanishi, 1940).

Distribution. China (Jilin), Korea, Japan.

Remarks. Larvae of *Epeorus latifolium* cannot be distinguished from those of *E. pellucidus*. Imanishi (1940) reported *E. latifolium* from Liaoning porvince, but it may belong to *E. pellucidus* judging from the distributional data.

46. *Epeorus pellucidus* (Brodsky) (Fig. 33)

Cinygma pellucida Brodsky, 1930: 35.

Diagnosis. Body light brown to brown. Frontal margin of head with light spots, with row of hairlike setae. Abdominal terga with paired dark markings. Outer margin of femora with row of long hairlike setae. Tails two.

Measurement (mm). ♂: BL: 13.2, HL: 2.2, HW: 3.2, AL: 2.1, EW: 1.1, DE: 0.6, CL: 10.2. ♀: BL: 15.2, HL: 2.6, HW: 3.7, AL 2.0, EW: 1.1, DE: 0.6, CL: 14.0.

Materials examined. Heilongjiang Heihe Xingfuxiang (4L, 26 VII 1997, YTQ), Sijiazi (7L, 26 VII 1997, YTQ), Hailinshi Pinghe (2L, 28 VII 1997 YTQ), Hailinshi Sanbuluo (20L, 28 VII 1997, YTQ), I; Hailinshi Shunqiao (30L, 28 VII 1997, YTQ), Shangzhiishi Maorshanzen (61L, 4 VI 1999, YTQ), Yanshouxian Yanhe (21L, 6 VI 1999, YTQ), Mudanjiangshi Xinfeng (24L, 29 VII 1997, YTQ), Mudanjiangshi Liuchun (86L, 29 VII 1997, YTQ), Ninganxian (Dongjingchengzhen (33L, 15 VII 2000, YTQ). Jilin Hailin Huanggang (5L, 18 VII 2000, YTQ), Yanjishi Xiaowudao (21L, 27 VII 2000, YTQ), Yanjishi Xiaowudao (2L, 30 VI 2000, YTQ), Yanjishi Yanxiaoqiao (4L, 2 VIII 2000, YTQ), Antu Toudaobaihe (103L, 3 VI 1997, YJB), Antu Baihe (10L, VI 1997, YJB), Antu Sandao (23L, VI 1997, YJB), Antu Sidaobaihe (25L, VI 1997, YJB), Wolong (3L, 6 VI 1997, YJB).

Distribution. China (Jilin), Korea, Japan.

47. *Heptagenia flava* Rostock

Heptagenia na: Imanishi, 1940: 256.

Heptagenia flava Rostock, 1878: 76.

Materials examined. Not available.

Locality. Heilongjiang Mudanjiangjingpohu (Imanishi, 1940).

Distribution. China (Heilongjiang, Neimenggu), Mongolia, Russia, Japan.

Remarks. Imanishi (1940) reported larvae of *Heptagenia* na from Heilongjiang province which was subsequently associated with *H. flava* (see Bae et al., 2000).

48. *Heptagenia kihada* Matsumura (Figs. 34, 90, 154, 172, 181, 202)

Heptagenia kihada Matsumura, 1931: 1478.

Diagnosis. Body brown. Lacinial region of left mandible with a single hair. Thorax brown with irregular light markings. Tibiae slightly shorter than femora. Gills on abdominal segments 1–7, apically pointed.

Measurement (mm). BL: 8.8, HL: 1.35, HW: 2.2, AL: 2.2, EL: 1.1, EW: 0.7, DE: 0.75, CL: 12.0, TF: 10.5.

Materials examined. Heilongjiang Ninganxian Dongjingchengzhen (1L, 15 VII 2000, YTQ).

Distribution. China (Heilongjiang), Japan, Mongolia, Russia.

49. *Heptagenia kyotoensis* Gose (Figs. 35, 48, 77, 91, 105, 133, 155, 173, 182)

Heptagenia kyotoensis Gose, 1963: 143.

Diagnosis. Body brown with light markings. Frontal margin of head without marking. Lateral pronotum and mesonotum with paired light stripes. Abdominal tergum with paired light longitudinal stripes, with paired markings. Abdominal segments 1–7 with tufts of filamentous gills. Caudal filaments very long.

Measurement (mm). BL: 11.1, AL: 1.9, CL: 17.5.

Materials examined. Heilongjiang Ninganxian Dongjingchengzhen (1L, 15 VII 2000, YTQ).

Distribution. China (Heilongjiang), Korea, Japan.

50. *Heptagenia sulphurea albicauda* Kluge (Fig. 36)

Heptagenia sulphurea albicauda Kluge, 1987: 318.

Diagnosis. Body reddish brown. Abdominal terga 3–10 with light paired markings; posterior margin of abdominal segment 1–10 with acute processes. Caudal filament with whorls of setae at end of each segment.

Measurement (mm). BL: 10, HL: 2.4, HW: 3.3, AL: 4, EL: 1.3, EW: 0.8, DE: 0.4, TF: 14.

Materials examined. Jilin Yanji (1L, 6 VI 1997, YJB).

Distribution. China (Jilin), Russia.

51. *Iron aesculus* (Imanishi) (Figs. 37, 49, 78, 93)

Epeorus aesculus Imanishi, 1934: 384.

Diagnosis. Body light brown to brown. Frontal margin of head lighter, with dark median stripe. Abdominal segments 2–7 with two lateral spines. Gill lamellae on abdominal segment 1 and 7 meeting beneath body. Tails two, without setae.

Measurement (mm). BL: 10.4, HL: 1.9, HW: 2.7, AL: 2.3, EL: 0.8, EW: 0.4, DE: 0.9, CL: 12.0.

Materials examined. Jilin Antu Toudaobaihe (1L, 3 VI 1997, YJB), Antu Eradaobaihe (19L,

3.VI 1997, YJB), Antu Baihe (14L, VI 1997, YJB), Antu Sandao (32L, VI 1997, YJB), Wolong Xianfeng (78L, 6 VI 1997, YJB), Helong Songxiaping (3L, VI 1997, YJB).

Distribution. China (Jilin), Korea, Russia.

Family Isonychiidae

52. *Isonychia japonica* (Ulmer) (Figs. 38, 79, 94, 106, 156, 174)

Chirotonetes japonicus Ulmer, 1919: 59.

Diagnosis. Body brown with light median longitudinal stripe. Forefemora and foretibiae with rows of hairlike setae. Abdominal segments 1–7 with both lamellate and filamentous gills.

Measurement (mm). BL: 15.8, AL: 5.0, EW: 1.4, DE: 0.3, CL: 8.1.

Materials examined. Jilin Yanjishi Xiaowudao (2L, 30 VI 2000, YTQ).

Distribution. China (Jilin), Korea, Mongolia, Russia, Japan.

53. *Isonychia na*

Isonychia na: Imanishi, 1940: 214.

Materials examined. Not available.

Locality. Heilongjiang Hailaer (Imanishi, 1940).

Distribution. China (Heilongjiang).

Remarks. Imanishi (1940) reported larva of this informal species from Heilongjiang province.

Family Leptophlebiidae

54. *Choroterpes altioculus* Kluge

Choroterpes trifurcata Uéno: Imanishi, 1940: 189.

Choroterpes (Euthraulus) altioculus Kluge, 1984: 722.

Materials examined. Not available.

Locality. Liaoning Dalian (Imanishi, 1940).

Distribution. China (Liaoning), Korea, Russia Japan.

Remarks. Bae *et al.* (1994) corrected Imanishi's (1940) report of the larvae of *Choroterpes trifurcata* Uéno from Japan and Korea as *C. altioculus* Kluge. Imanishi's Manchurian materials also may belong to *C. altioculus*.

55. *Paraleptophlebia chocolata* Imanishi (Figs. 39, 81, 107, 134, 135, 157, 188, 203)

Paraleptophlebia chocolata Imanishi, 1937b: 330.

Diagnosis. Body chocolate brown. Head with three light round spots. Gills 1–7 bi-forked; tracheae of the gills with small lateral branches.

Measurement (mm). ♂: BL: 8.0, HW: 1.5, HL: 1.3, EW: 0.4, EL: 0.72, DE: 0.35, CL: 5.0.

♀: BL: 9.4, HW: 1.3, HL: 1.35, EW: 0.3, EL: 0.4, DE: 0.85.

Materials examined. Heilongjiang Hailinshi Sanbuluo (2L, 28 VII 1997, YTQ), Shangzhishi Maorshanzhen (3L, 4 VI 1999, YTQ).

Distribution. China (Heilongjiang), Korea, Mongolia, Russia.

56. *Thraulus na*

Thraulus na: Imanishi, 1940: 189.

Materials examined. Not available.

Locality. **Liaoning** Erdaohezi (Imanishi, 1940).

Distribution. China (Liaoning).

Remarks. Imanishi (1940) reported larva of this undetermined species from Liaoning province.

57. *Thraulus YUa* (Figs. 40, 50, 80, 95, 108, 158, 175, 204, 205)

Diagnosis. Body grayish brown. Anterior margin of pronotum with hairlike setae. Mesonotum with irregular markings. Abdominal segment 7–9 with distinct posterolateral processes. Gills 1 single; gills 2–7 multi-forked.

Measurement (mm). BL: 6.2, HL: 1.4, HW: 1.35, EL: 0.65, EW: 0.4, DE: 0.25, CL: 9.0.

Materials examined. **Heilongjiang** Mudanjiangshi Dongjingcheng (1L, 15 VII 2000, YTQ).

Distribution. China (Heilongjiang).

Remark. This species is distinguished from *Thraulus na* by possessing relatively larger abdominal gills and well developed posterolateral projections on abdominal segment 7.

Family Neoephemeridae

58. *Potamanthellus chinensis* Hsu

Potamanthellus chinensis Hsu, 1935: 321.

Materials examined. Not available.

Locality. **Jilin** Beishan (You and Gui, 1995).

Distribution. China (Jilin), Korea.

Remarks. Hsu (1935) described adult of this species from Jilin province.

Family Polymitarcyidae

59. *Ephoron shigae* (Takahashi) (Figs. 41, 51, 98, 99, 136, 206)

Polymitarcis shigae Takahashi, 1924: 379.

Diagnosis: Body light yellow to light brown. Head with round clypeal process. Mandibles developed into long tusks with small tubercles.

Measurement (mm). BL: 5.2, HL: 0.2, HW: 0.3, EL: 0.2, EW: 0.3, DE: 1.25, CL: 1.5, TF: 1.9.

Materials examined. **Heilongjiang** Shangzhishi Maorshanzhen (2L, 4 VI 1999, YTQ), Hailinski Pinghe (4L, 28 VII 1997, YTQ).

Distribution. China (Heilongjiang), Japan, Korea, Russia.

60. *Ephoron virgo* (Oliver)

Ephoron virgo Olivier, 1791: 404.

Materials examined. Not available.

Locality. **Jilin** (You and Gui, 1995).

Distribution. China (Jilin), Russia.

Remarks. You and Gui (1995) reported adult of this species from Jilin province.

Family Potamanthidae

61. *Potamanthus luteus oriens* Bae and McCafferty (Figs. 42, 61, 207)

Potamanthus (Potamanthus) luteus oriens Bae and McCafferty, 1991: 54.

Diagnosis. Body brown with light markings, dorso-ventrally flattened. Mature male compound eyes relatively large. Mandibles 0.3 x length of head. Forefemora with dorsal transverse row of 10–15 simple-shout setae. Gills 2–7 each with 20–30 marginal fibrillae arranged in single row in each lamella.

Measurement (mm). BL: 11.5, HL: 1.7, HW: 2.6, AL: 2.4, EL: 1.7, EW: 0.8, DE: 1.0, CL: 6.8.

Materials examined. Heilongjiang Ninganxian Dongjingchengzhen (2L, 15 VII 2000, YTQ).

Distribution. China (Heilongjiang), Korea, Japan.

62. *Rhoenanthus youi* (Wu)

Neopotamanthus youi Wu, 1986: 401.

Materials examined. Not available.

Locality. Jilin Antuxian, Fusongxian (You and Gui, 1995).

Distribution. China (Jilin).

Remarks. Based on distributional data of the species of Potamanthidae (Bae and McCafferty, 1991), this species probably belongs to *Rhoenanthus coreanus* (Yoon and Bae) which is distributed in the Korean peninsula and Russian Far East. *R. youi* occurs in southern China.

Family Siphlonuridae

63. *Siphlonurus binotatus* (Eaton)

Siphlurus binotatus Eaton, 1892: 302.

Materials examined. Not available.

Locality. Heilongjiang Heihenkuanhe (Imanishi, 1940).

Distribution. China (Heilongjiang), Japan.

64. *Siphlonurus chankae* Tshernona (Fig. 43)

Siphlonurus chankae Tshernona, 1952: 263.

Diagnosis. Body yellowish brown. Wingpads without dark spots. Legs with single dark transverse band on femora and tibiae of all legs. Posterolateral spines of abdomen poorly developed. Gills 1 posterior margin of dorsal and ventral lamellae slightly notched.

Measurement (mm). BL: 15.4, HL: 2.5, HW: 2.0, EL: 1.2, EW: 0.8, DE: 0.5, CL: 6.0, TF: 5.7.

Materials examined. Heilongjiang Heihe Xingfuxiang (76L, 26 VII 1997, YTQ), Sijiazi (129L, 26 VII 1997, YTQ), Hailinshi Pinghe (6L, 28 VII 1997, YTQ), Hulan (3L, 23 VII 1999, YTQ), Yanjishi Xiaowudao (6L, 30 VI 2000, YTQ).

Distribution. China (Heilongjiang), Korea.

65. *Siphlonurus immanis* Kluge (Figs 44, 55, 96, 137, 149, 176)

Siphlonurus immanis Kluge 1985: 18.

Diagnosis. Body color grayish brown. Wingpads without dark spots. Legs with single dark transverse band on femora and tibiae of all legs. Posterolateral spines of abdomen well developed. Gills 1 posterior margin of dorsal and ventral lamellae deeply notched.

Measurement (mm). ♂: BL: 15.5, HL: 3.0, HW: 2.4, AL: 2.1, EL: 1.4, EW: 1.0, DE: 1.0, CL: 8.0, TF: 7.0, ♀: BL: 13.5, AL 1.3, CL: 6.8, TF: 6.2.

Materials examined. Jilin Yanjishi Xiaowudao (2L, 27 VII 2000, YTQ).

Distribution. China (Jilin).

Acknowledgements This work was supported by the grant No. R01-2001-000-00086-0 from the Basic Research Program of the Korea Science & Engineering Foundation.

REFERENCES

- Bae, Y.J. 1995. *Ephemerella separigata*, a new species of Ephemeroptera (Insecta: Ephemeroptera) from Korea. Korean J. Syst. Zool. 11: 159–166.
- Bae, Y.J. 1997. A historical review of Ephemeroptera systematics in Northeast Asia. In Landolt, P. and M. Sartori. (eds). *Ephemeroptera and Plecoptera*. MTL, Fribourg, Switzerland. pp. 405–417.
- Bae, Y.J. and G.C. Liu. 1999. Mayflies (Ephemeroptera) from Changbaishan area in Northeast Asia. Entomol. Res. Bull. (KEI). 25: 1–6.
- Bae, Y.J. and W.P. McCafferty. 1991. Phylogenetic systematics of the Potamanthidae (Ephemeroptera). Trans. Am. Entomol. Soc. 117: 1–143.
- Bae, Y.J., J.E. Lee, and I.B. Yoon. 2000. Northeast Asian Ephemeroptera in Imanishi's 1940 report. Entomol. Sci. 3: 391–397.
- Bajkova, O.Y. 1962. New species of the genus *Ephemerella* Walsh (Ephemeroptera, Ephemerellidae) from mountain tributaries of the Amur basin. Izv. Tikhookean. Nauchn. Issled. Inst. Rybn. Khoz. Okeanogr. 48: 202–205.
- Bajkova, O.Y. 1967. New species of *Ephemerella* Walsh (Ephemeroptera, Ephemerellidae) from Far East and East Siberia. Entomol. Obozr. 46: 327–335.
- Bajkova, O.Y. 1980. A new species of mayfly (Ephemeroptera, Ephemerellidae) from Primorye region. Entomol. Obozr. 79: 74–76.
- Bengtsson, S. 1909. Beitrage zur Kenntnis der palaarktischen Ephemereniden. Lunds Univ. Arsbok, N. F., Afd. 2, 5(4): 1–19.
- Broadsky, K.A. 1930. Contributions to the fauna of Ephemeroptera of Southern Siberia. Entomol. Obozr. 24: 31–40.
- Eaton, A.E. 1870. On some new British species of Ephemeroidea. Trans. Entomol. Soc. London 18: 1–8.
- Eaton, A.E. 1892. On two new and some other Japanese species of Ephemeroidea. Entomol. Mon. Mag. 38: 302–303.
- Gose, K. 1963. Two new mayflies from Japan. Kontyu 31: 142–145.
- Gui, H. 1985. A catalogue of the Ephemeroptera of China. Bull. Nanjing Normal Univ. (Nat. Sci.) 4: 79–97.
- Hsu, Y.C. 1931–32. Two new species of mayflies from China (Order Ephemeroidea). Peking Nat. Hist. Bull. 6: 39–41.
- Hsu, Y.C. 1935–36. New Chinese mayflies from Kiangsi province (Ephemeroptera). Peking Nat. Hist. Bull. 10: 319–326.
- Imanishi, K. 1930. Mayflies from Japanese torrents. I. New mayflies of the genera *Acentrella* and *Ameletus*. Trans. Nat. Hist. Soc. Formosa 20: 263–267.
- Imanishi, K. 1934. Mayflies from Japanese torrents. IV. Notes on the genera *Epeorus*. Annot. Zool. Jap. 14: 381–395.
- Imanishi, K. 1935. Mayflies from Japanese torrents. V. Notes on the genera *Cinygma* and *Heptagenia*. Annot.

- Zool. Japon. 15: 213-223.
- Imanishi, K. 1936. Mayflies from Japanese torrents. VI. Notes on the genera *Ecdyonurus* and *Rhithrogena*. Annot. Zool. Jap. 15: 538-549.
- Imanishi, K. 1937a. Mayflies from Japanese torrents. VII. Notes on the genus *Ephemerella*. Annot. Zool. Jap. 16: 321-329.
- Imanishi, K. 1937b. Mayflies from Japanese torrents. VIII. Notes on the genera *Paraleptophlebia* and *Baetis*. Annot. Zool. Jap. 16: 330-339.
- Imanishi, K. 1940. Ephemeroptera of Manchoukuo, inner Mongolia, and Chosen. In Kawamura, T. (ed.), Report of the Limnological Survey of Kwantung and Manchoukuo. Pp. 169-263.
- Kazlauskas, R. 1963. New and little-known mayflies (Ephemeroptera) in the fauna of the USSR. Entomol. Obozr. 42: 582-589.
- Kluge, N.J. 1983a. New and little known mayflies of the family Baetidae (Ephemeroptera) from the Primorye. Entomol. Rev. 83: 53-68.
- Kluge, N.J. 1983b. New and little-known mayflies of Far-East USSR. Genus *Ecdyonurus* Etn. (Ephemeroptera, Heptageniidae). Ekologiya i Cistematika Presnovodnykh Organizmov Dalnego Vostoka. Pp. 27-36.
- Kluge, N.J. 1984. Mayflies of the subgenus *Euthraulus* Barn. (Ephemeroptera, Leptophlebiidae, genus *Choroterpes*) of the fauna of the USSR. Entomol. Obozr. 63: 722-728.
- Kluge, N.J. 1985. On Far Eastern mayflies of the group *aestivalis* of the genus *Siphlonurus* Etn. (Ephemeroptera, Siphlonuridae). Vestnik Lenigrad. Univ., 10: 12-20.
- Kluge, N.J. 1987. Mayflies of the genus *Heptagenia* Walsh (Ephemeroptera, Heptageniidae) of the fauna of the USSR. Entomol. Obozr. 66: 302-320.
- Lee, J.W., Y.T. Quan and J.C. Jung. 2001. Mayflies from Heilongjiang and Jilin provinces in Northeastern China. Korean J. Entomol. 31: 15-23.
- Linnaeus, C. 1761. Fauna Suecica. Ed. II. stockholmae. No. 1477.
- Malzacher, P. 1996. *Caenis nishinoae*, a new species of the family Caenidae from Japan (Insecta: Ephemeroptera). Stutt. Beitr. Natur. Ser. A (Biol.) 547: 1-5.
- Matsumura, S. 1911. Erster Beitrag zur Insekten Fauna von Sachalin. J. Coll. Agr., Tohoku Imp. Univ., Sapporo. 4: 6-7.
- Matsumura, S. 1931. 6000 Illustrated Insects of the Japanese Empire. Pp. 1465-1480.
- McLachlan, R. 1875. A sketch of our present knowledge of the neuropterous fauna of Japan. Trans. Entomol. Soc. London. Pp. 167-190.
- Navas, L. 1912. Quelques Nevropteres de la Siberie meridionale-orientale. Revue Russe d' Entomol. 12: 414-422.
- Olivier, G.A. 1791. Artikel "Ephemerida". Encyclopedie Methodique. 6: 404-422.
- Pactl, J. 1994. *Ephacerella*, a replacement name for *Acerella* Allen, 1971 (Ephemeroptera), nec Berlese, 1909 (Protura). Ent. News 105: 283-284.
- Rostock, M. 1878. Die Ephemeren und Psociden Sachsen mit Berücksichtigung der meisten übrigen deutschen Arten. Jber. Ver. Naturk. Zwickau. 1878: 76-100.
- Su, C.R. and D.S. You. 1988. Two new species of genus *Ephemerella* subgenus *Ephemerella* (Ephemeroptera: Ephemellidae). J. Nanjing Normal Univ. (Nat. Sci.). 3: 63-67.
- Su, C.R. and D.S. You. 1989. A new species of genus *Ephemerella* subgenus *Ephemerella* (Ephemeroptera: Ephemellidae). Acta Zootaxon. Sinica. 14: 181-185.
- Su, C.R. and H. Gui. 1995. The first record of the genus *Drunella* in China with description of a new species (Ephemeroptera: Ephemellidae). Acta Zootaxon. Sinica. 20: 451-454.
- Takahashi, Y. 1924. New species of mayflies from Japan. J. Zool. 63: 377-380.
- Tiunova, T.M. 1990. Descriptions of a new species of mayflies of the genus *Cinygmulia* McD. (Ephemeroptera, Heptageniidae) and the larva of *Rhoenanthus rohdendorfi* Tshern. (Potamanthidae) from Primorye Territory. Entomol. Obozr. 69: 814-819.
- Tshernova, O.A. 1949. Mayfly nymphs from Lake Teletskoye and River Bija. Trudi Zool. Inst. Akad. Nauk. 7: 139-158.
- Tshernova, O.A. 1952. Mayflies of the Amur River Basin and nearby waters and their role in the nutrition of Amur fishes. Trudi Amurskoi Ikhtiologicheskoi Ekspeditsii. 1945-1949. 3: 229-360.
- Tshernova, O.A. and V.V. Belov. 1982. Systematic position and synonymy of *Cinygma tibiale* Ulmer 1920 (Ephemeroptera, Heptageniidae). Ent. Mitt. aus dem Hamb. Zool. Mus. und Ins. 7: 193-194.

- Uéno, M. 1928. Some Japanese mayfly nymphs. Mem. Coll. Sci., Kyoto Imp. Univ., Kyoto 4: 19-63.
- Ulmer, G. 1919. Neue Ephemeropteren. Arch. Naturg. 85: 1-80.
- Wu, X.Y. and D.S. You. 1986. A new genus and species of Potamanthidae from China (Ephemeroptera). Acta Zootaxon. Sinica. 11: 401-405.
- You, D.S. and H. Gui. 1995. Economic Insect Fauna of China. Fasc. 48. (Ephemeroptera). Science Press. Beijing, China.

(Received: June 15, 2002, Accepted: July 8, 2002)

EXPLANATIONS OF FIGURES

- Figs. 1–44.** Larval body, dorsal: 1. *Ameletus costalis*; 2. *A. montanus*; 3. *Acentrella sibirica*; 4. *Baetis fuscatus*; 5. *B. pseudothermicus*; 6. *Baetiella tuberculata*; 7. *Cloeon dipterum*; 8. *Procloeon pennulum*; 9. *Brachycercus YUa*; 10. *Caenis na*; 11. *Cincticostella levanidovae*; 12. *C. tshernovae*; 13. *Drunella cryptomeriae*; 14. *D. lepnevae*; 15. *D. solida*; 16. *Ephemerella aurivillii*; 17. *E. dentata*; 18. *E. kozhovi*; 19. *Serratella setigera*; 20. *S. zapekinae*; 21. *Uracanthella rufa*; 22. *Ephemerella orientalis*; 23. *E. separata*; 24. *E. strigata*; 25. *Cinygmulia brunnea*; 26. *C. hirasana*; 27. *C. kurenzovi*; 28. *C. levanidovi*; 29. *Ecdyonurus dracon*; 30. *E. kibunensis*; 31. *E. levis*; 32. *Epeorus curvatus*; 33. *E. pellucidus*; 34. *Heptagenia kihada*; 35. *H. kyotoensis*; 36. *H. sulphurea albicauda*; 37. *Iron aesculus*; 38. *Isonychia japonica*; 39. *Paraleptophlebia chocolata*; 40. *Thraulus YUa*; 41. *Ephoron shigae*; 42. *Potamanthus luteus oriens*; 43. *Siphlonurus chankae*; 44. *S. immanis*. (scale = 1 mm).

Figs. 45–51. Head, dorsal: 45. *Drunella solida*; 46. *Cinygmulia hirasana*; 47. *Ecdyonurus kibunensis*; 48. *Heptagenia kyotoensis*; 49. *Iron aesculus*; 50. *Thraulus YUa*; 51. *Ephoron shigae*.

Figs. 52–54. Thorax, dorsal: 52. *Cincticostella levanidovae*; 53. *Ephemerella dentata*; 54. *Cinygmulia hirasana*.

Figs. 55–58. Abdomen, dorsal: 55. *Siphlonurus immanis*; 56. *Baetis pseudothermicus*, 3rd–6th; 57. *Baetis pseudothermicus*, posterior margin; 58. *Cloeon dipterum*, 10th posterior margin.

Fig. 59. Thorax, ventral: *Cincticostella tshernovae*.

Figs. 60–61. Caudal filaments: 60. *Procloeon pennulum*, basal; 61. *Potamanthus luteus oriens*, middle segments.

Figs. 62–82. Legs: 62. *C. levanidovae* hindleg; 63. *Cincticostella levanidovae*, foreleg; 64. *Procloeon pennulum*, hindleg; 65. *C. tshernovae*, foreleg; 66. *C. tshernovae*, midleg; 67. *Drunella lepnevae*, foreleg; 68. *D. solida*, foreleg; 69. *Ephemerella dentata*, foreleg; 70. *Serratella zapekinae*, foreleg; 71. *Uracanthella rufa*, foreleg; 72. *Ephemerella strigata*, foreleg; 73. *Cinygmulia hirasana*, foreleg; 74. *Ecdyonurus dracon*, foreleg; 75. *E. kibunensi*, foreleg; 76. *Epeorus curvatus*, foreleg; 77. *Heptagenia kyotoensis*, foreleg; 78. *Iron aesculus*, foreleg; 79. *Isonychia japonica*, foreleg; 80. *Thraulus YUa*, foreleg; 81. *Paraleptophlebia chocolata*, foreleg; 82. *Acentrella sibirica*, foreleg.

Figs. 83–99. Labrum, dorsal: 83. *Ameletus costalis*; 84. *Baetis pseudothermicus*; 85. *Procloeon pennulum*; 86. *Cincticostella levanidovae*; 87. *Ephemerella dentata*; 88. *Serratella zapekinae*; 89. *Ecdyonurus dracon*, ventral; 90. *Heptagenia kihada*, ventral; 91. *H. kyotoensis*, ventral; 92. *Epeorus curvatus*; 93. *Iron aesculus*, left-dorsal, right-ventral; 94. *Isonychia japonica*; 95. *Thraulus YUa*, left-dorsal, right-ventral; 96. *Siphlonurus immanis*, left-dorsal, right-ventral; 97. *Acentrella sibirica*; 98. *Ephoron shigae*; 99. *E. shigae*, transverse section, middle part of labrum.

Figs. 100–108. Hypopharynx, ventral: 100. *Drunella solida*; 101. *Ephemerella dentata*; 102. *Uracanthella rufa*; 103. *Ecdyonurus dracon*; 104. *Epeorus curvatus*; 105. *Heptagenia kyotoensis*; 106. *Isonychia japonica*; 107. *Paraleptophlebia chocolata*; 108. *Thraulus YUa*.

Figs. 109–139. Mandibles, ventral: 109. *Ameletus costalis*, right; 110. *A. montanus*, right; 111. *Baetis pseudothermicus*, right; 112. *B. pseudothermicus*, left; 113. *Procloeon pennulum*, right; 114. *Brachycercus YUa*, right; 115. *Cincticostella levanidovae*, right; 116. *C. levanidovae*, left; 117. *C. tshernovae*, left; 118. *Drunella solida*, left; 119. *D. cryptomeriae*, right; 120. *Ephemerella kozhovi*, left; 121. *Drunella lepnevae*, right; 122. *D. lepnevae*, left; 123. *Serratella zapekinae*, right; 124. *Uracanthella rufa*; 125. *U. rufa*, left; 126. *Acentrella sibirica*, right; 127. *A. sibirica*, left; 128. *Baetis fuscatus*, right; 129. *B. fuscatus*, left; 130. *Ecdyonurus dracon*, right; 131. *E. dracon*, left; 132. *E. levis*, right; 133. *Heptagenia kyotoensis*, right; 134. *Ecdyonurus strigata*, right; 135. *P. chocolata*, left; 136. *Ephoron shigae*, right; 137. *Siphlonurus immanis*, right; 138. *Caenis na*, right; 139. *Caenis na*, left.

Figs. 140–161. Maxilla, ventral: 140. *Ameletus costalis*; 141. *Brachycercus YUa*; 142. *Cincticostella levanidovae*; 143. *C. tshernovae*; 144. *Drunella lepnevae*; 145. *D. solida*; 146. *Ephemerella kozhovi*; 147. *Uracanthella rufa*; 148. *Ephemerella strigata*; 149. *Siphlonurus immanis*; 150. *E. dracon*; 151. *Ecdyonurus kibunensis*; 152. *E. levis*; 153. *Epeorus curvatus*; 154. *Heptagenia kihada*; 155. *H. kyotoensis*; 156. *Isonychia japonica*; 157. *Paraleptophlebia chocolata*; 158. *Thraulus YUa*; 159. *Acentrella sibirica*; 160. *Baetis fuscatus*; 161. *Caenis na*.

Figs. 162–177. Labium: 162. *Procloeon pennulatum*, left-ventral, right-dorsal; 163. *Cincticostella levanidovae*, right-dorsal, left-ventral; 164. *C. tshernovae*, ventral; 165. *Drunella solida*, ventral; 166. *Ephemerella kozhovi*, ventral; 167. *Uracanthella rufa*, ventral; 168. *Ephemera strigata*, ventral; 169. *Ecdyonurus dracon*, ventral; 170. *E. kibunensis*, ventral; 171. *E. levis*, ventral; 172. *Heptagenia kihada*, ventral; 173. *H. kyotoensis*, ventral; 174. *Isonychia japonica*, left-ventral, right-dorsal; 175. *Thraulus YUa*, ventral; 176. *Siphlonurus immanis*, ventral; 177. *Baetis fuscatus*, ventral.

Figs. 178–182. Labial palpus: 178. *Baetis pseudothermicus*; 179. *Ecdyonurus dracon*; 180. *E. levis*; 181. *Heptagenia kihada*; 182. *H. kyotoensis*.

Figs. 183–188. Tarsal claw: 183. *Cloeon dipterum*, hindclaw; 184. *Brachycercus YUa*, hindclaw; 185. *Cincticostella tshernovae*, foreclaw; 186. *Serratella zapekinae*, foreclaw; 187. *Uracanthella rufa*, hindclaw; 188. *Paraleptophlebia chocolata*, foreclaw.

Figs. 189–209. Gills: 189. *Ameletus costalis*, 3rd; 190. *Brachycercus YUa*, 2nd; 191. *Caenis na*, 2nd; 192. *Cincticostella tshernovae*, 1st; 193. *C. tshernovae*, 4th; 194. *Drunella cryptomeria*, 1st; 195. *D. solida*, 1st; 196. *D. solida*, 5th; 197. *Ephemerella dentata*, 1st; 198. *E. kozhovi*, 1st; 199. *E. kozhovi*, 5th; 200. *E. kozhovi*, 2nd; 201. *Serratella zapekinae*, 1st; 202. *Heptagenia kihada*, 3rd; 203. *Paraleptophlebia chocolata*, 3rd; 204. *Thraulus YUa*, 1st; 205. *Thraurus YUa*, 2nd.; 206. *Ephoron shigae*, 3rd; 207. *Potamanthus luteus oriens*, 3rd; 208. *Procloeon pennulatum*, 1st; 209. *Ephemera strigata*, 1st.













