

日本產蜉蝣新種

(大正十三年八月十日受領)

高橋 雄一

此記事を草するに當り岸田氏より多大の御教導にあつかり、又丸毛博士及木下理學士より色々御指導下されし處を此處に記入して深謝す。

I. アメフリカゲロウ (新稱) *Clivon maihonis*, n. sp. (Fig. 3)

雄はアオバカゲロウに非常によく似て居るが雌は翅に少しづつ色がついて居ないので直ちに彼と區別される。翅の外縁にある間投脈が一對であるのに、本属の他の多くのものと區別される。鉗子はフタバカゲロウに似て居る。

Female (in alcohol)

Eyes black; head and thorax rufous-fuscous. Abdomen in segment 2-7 testaceous, with many marks or spots which are in shape oblong and are rufous-fuscous of colour, and some of them with outline black; in segment 8 and 9 rufous-fuscous; remain testaceous, also caudal setae and all legs. Wings vitreous, contained two intercarys at the margin, as *Battis* or *Clivon fergusoni*, PETERSON.

Length of body: 4 mm.

Length of wing: 4.5 mm.

Loc: Shiga: Nagano.

Time of appearance: August.

II. アオバカゲロウ *Clivon Okumoti*, n. sp. (Fig. 1 and 2)

本種は岡本博士が岡崎氏の爲に新種として査定せし *Clivon semonis*, OKAMOTO なるを思ひしも記載なき爲同定するを得ず、こゝに新種として擧げたり。

雌の前翅の前縁が綠色なる故容易に前記の種類及フタバカゲロウを區別される。大抵はフタバカゲロウとアメフリカゲロウの中間にあり。

鉗子はアメフリカゲロウに極めてよく似たり。

Male (in alcohol)

Turbinate eyes reddish orange, basal eyes black. Head and thorax brownish yellow; legs tinted yellow slightly; wings vitreous. Abdominal segment 2-7 translucent : segment 8-10 cretaceous ; and these dorsum red, except segment 10.

Forces translucent, but basal segment cretaceous. Caudal setae colourless, with joinings red.

Length of body : 4.5 mm.




Fig. 1. Wing of *Cleon Okamotoi*, ♂.
 " 2. Forcep of *Cleon Okamotoi*, ♂.
 " 3. Wing of *Cleon miyakonis*, ♂.
 " 4. Penis and Forcep of *Ecdyurus yoshidai*, ♂.
 " 5. Penis and Forcep of *Ecterus ikmonis*, ♂.
 " 6. Part of Forewing of *Polyuktarsis shigae*, ♂.

Length of caudal seta : 11 mm.

Loc. : Tokyo; Kanazawa; Shizuoka; Okayama.

Time of appearance : July to September.

雌雄々々相似たり。雌は複眼黑色、體及脚白色。口下部の脛端、縫端、附節及胸、腹脊は赤褐色、尾毛は白色な

Male (dried)

Eyes black; head and prothorax greyish testaceous, meta-and mesothorax fuscous, Abdomen testaceous, caudal setae white, with joinings grey. In the fore legs femur testaceous, the upper side of terminal edge reddish black; tibiae testaceous, with many spots reddish black, meso-and metathoracic legs testaceous.

Length of body : 9 mm.

Length of forewing : 9 mm.

Loc. : Shiga, Shizuoka.

Time of appearance: August and September.

■ ヤカノカケロウ (新種) *Flynnius ikunoris*, n. sp. (Fig. 5)

雄、複眼灰色、頭部黑色、前脚褐色、腹端部の後縁に各々黒褐色の横紋あり。翅は墨無色にして脈は褐色な

Male (in alcohol)

Eyes cinereous pale. Pronotum picaceous, each side reddish fuscous. Lateral side of prothorax has a dot yellow; and mesonotum with a picaceous streak in the center and in each side with a oblong dots testaceous. Wings, vitreous, tinted dark yellow slightly, and the pterostigmata space brownish black. All legs picaceous. Abdominal segment testaceous, with picaceous joinings, lateral suture yellow; Forceps and caudal setae picaceous.

Length of body, 13.5 mm.

Length of forewing: 14 mm.

Length of post wing : 5 mm.
Length of caudal seta : 37 mm.

Loc. Shauki.

Time of appearance : March.

ヌマヒラタカロウ (新種) *Ridunus yoshidae*, n. sp. (Fig. 4)

若、雄はヘビイロアゲハ、複眼は雄は大で、雌は比較的小で、前脚は灰色だ。翅は黒色で、翅の前縁は緑色、體は緑黄色。雄の腹端は半透明、翅の脇腹脊は褐色だ。

日本産
昆蟲
新種

Female (living)
Compound eyes intermediate darkish brown, and is marked at the tip two short red dash ; at the side greenish yellow, with three rings piceous.
Ocelli colourless, the bottom is in colour black. Antenna apical segment grey, subtranslucent, the other two segments deep brown.

Head greenish yellow, vertex darkish brown. Prothorax greenish yellow, notum narrow tinted darkish brown : mesothorax greenish yellow, notum darkish brown.
Abdomen in the segment 1-9 greenish yellow, and dorsum tinted brown, and the lateral sides have twig shaped marks which are in colour black, and ventral more greenish yellow of colour : of every segments with narrow black band at the segment. Last abdominal segment greenish yellow.

Tarsus and claws grey, femer yellowish grey, the other parts of legs greenish yellow. Wings vitreous, costal and radii us space tinted green. Caudal setae greenish grey, without colour rings.

Length of body : 10 mm.

Length of fore wing : 11 mm.

Length of post wing : 4 mm.

Length of caudal setae : 19 mm.

Loc. Shauki, Shiga, Fukushima.

Time of appearance, June to August.