

Michael Hubbard

MIRJANA TANASIEVIC

NEW FIND OF THE SPECIES
PARALEPTOPHLEBIA RUFFOI Biancheri, 1956
(Ephemeroptera, Leptophlebiidae)

NOV NALAZ VRSTE
PARALEPTOPHLEBIA RUFFOI Biancheri, 1956
(Ephemeroptera, Leptophlebiidae)

SEPARAT IZ »GODIŠNJAKA BIOLOŠKOG INSTITUTA
UNIVERZITETA U SARAJEVU«

VOL. 34 — 1981. GOD.

MIRJANA TANASIEVIC

Prirodno-matematicki fakultet, Sarajevo

NEW FIND OF THE SPECIES
PARALEPTOPHLEBIA RUFFOI Biancheri, 1956
(Ephemeroptera, Leptophlebiidae)

NOV NALAZ VRSTE
PARALEPTOPHLEBIA RUFFOI Biancheri, 1956
(Ephemeroptera, Leptophlebiidae)

The species *Paraleptophlebia ruffoi* was described according to male and female samples collected during the month of July on Monta Sibillina (Italy, Umbria) at the height of 1323 m above sea level (Biancheri, 1956). An other find of the same species, imago and subimago was mentioned by Grandi (1962) on the basis of material collected in the lake Trasimeno during April and May. According to these data Puthz (1978) considers *Paraleptophlebia ruffoi* as the endemic species of the central part of the Appenines, which Illies (1967), included in the region 3.

Belfiore and Giangrande (1979) have described nymphs of this species, on the basis of the material collected at the height of 350 m above sea level nearby the border between Umbria and Toscana, and in the same time, they have published new find of imagos in the lake Canterno (Lazio, Fiuggi), 538 m above sea. The imagos were collected in the middle of the June 1978.

All that indicates that this species of the central part of the Appenines emerges at the altitude of 350 m to 1323 m from early spring until the beginning of the summer.

In the material collected by Professor Mara Marinković-Gspodnetić from the running waters of Macedonia in June 1973, the species is recorded at one locality. It has been found in a small stream on the mountain Babuna, on the way Titov Veles — Prilep, at the altitude about 500 m, 3. 6. 1973, 3 ♂♂, 2 ♂♂ (s), 3 ♀♀.

Thus the area of the species *Paraleptophlebia ruffoi* is expanded in the South part of the Balkan peninsula, i.e. to the region 6. In other parts of the peninsula this species has not yet been known.

I would like to thank very much to Professor Mara Marinković-Gospodnetić for the material granted.

REZIME

Vrsta *Paraleptophlebia ruffoi* Bancheri poznata do sada iz centralnog dijela Apeninskog gorja nađena je na jednom lokalitetu u Makedoniji. Ova vrsta je nađena na malom livadskom potoku na planini Babuni, put Titov Veles — Prilep na n.v. oko 500 m. Time je areal vrste *Paraleptophlebia ruffoi* proširen na južni dio Balkanskog poluostrva odnosno na regiju 6 po Illies-u (1967).

LITERATURA

- Belfiore, C. et A. Giangrande, 1979. Descrizione della ninfa di *Paraleptophlebia ruffoi* Bancheri, 1956 (Ephemeroptera, Leptophlebiidae). *Fragm. Entomol.*, Roma, 15 (1): 53—58.
- Biancheri, E. 1956. Una nuova specie della famiglia Leptophlebiidae, *Paraleptophlebia ruffoi* n. sp. *Mem. Mus. Civ. St. Nat. Verona*, 5:191—194.
- Grandi, M. 1960. Ephemeroidea. *Fauna d'Italia*, Calderini, Bologna: 1—474.
- Grandi, M. 1962. Nota su alcuni Efemeroidei del Lago Trasimeno. *Riv. di Idrobiol.*, 1, (2—3): 179—187.
- Illies, J. 1967. Limnofauna Europaea. Eine Zusammenstellung aller die europäischen Binnengewässer bewohnenden mehrzelligen Tierarten mit Angaben über ihre Verbreitung und Ökologie. Fischer, Stuttgart, XV + +474pp., lpl.
- Putthz, V. 1978. Ephemeroptera in J. Illies, *Limnofauna Europaea*, Fischer, Stuttgart: XVII+532pp., lpl.